PROCEEDINGS OF SPIE

SPIDigitalLibrary.org/conference-proceedings-of-spie

Front Matter: Volume 8174

, "Front Matter: Volume 8174," Proc. SPIE 8174, Remote Sensing for Agriculture, Ecosystems, and Hydrology XIII, 817401 (20 October 2011); doi: 10.1117/12.913083

Event: SPIE Remote Sensing, 2011, Prague, Czech Republic

PROCEEDINGS OF SPIE

Remote Sensing for Agriculture, Ecosystems, and Hydrology XIII

**Christopher M. U. Neale
Antonino Maltese
Katja Richter**
Editors

**19–21 September 2011
Prague, Czech Republic**

Sponsored by
SPIE

Cooperating Organisations
EOS— European Optical Society
Remote Sensing and Photogrammetry Society (United Kingdom)

Published by
SPIE

Volume 8174

Proceedings of SPIE, 0277-786X, v. 8174

SPIE is an international society advancing an interdisciplinary approach to the science and application of light.

The papers included in this volume were part of the technical conference cited on the cover and title page. Papers were selected and subject to review by the editors and conference program committee. Some conference presentations may not be available for publication. The papers published in these proceedings reflect the work and thoughts of the authors and are published herein as submitted. The publisher is not responsible for the validity of the information or for any outcomes resulting from reliance thereon.

Please use the following format to cite material from this book:

Author(s), "Title of Paper," in *Remote Sensing for Agriculture, Ecosystems, and Hydrology XIII*, edited by Christopher M. U. Neale, Antonino Maltese, Katja Richter, Proceedings of SPIE Vol. 8174 (SPIE, Bellingham, WA, 2011) Article CID Number.

ISSN 0277-786X

ISBN 9780819488015

Published by

SPIE

P.O. Box 10, Bellingham, Washington 98227-0010 USA

Telephone +1 360 676 3290 (Pacific Time) · Fax +1 360 647 1445

SPIE.org

Copyright © 2011, Society of Photo-Optical Instrumentation Engineers

Copying of material in this book for internal or personal use, or for the internal or personal use of specific clients, beyond the fair use provisions granted by the U.S. Copyright Law is authorized by SPIE subject to payment of copying fees. The Transactional Reporting Service base fee for this volume is \$18.00 per article (or portion thereof), which should be paid directly to the Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923. Payment may also be made electronically through CCC Online at copyright.com. Other copying for republication, resale, advertising or promotion, or any form of systematic or multiple reproduction of any material in this book is prohibited except with permission in writing from the publisher. The CCC fee code is 0277-786X/11/\$18.00.

Printed in the United States of America.

Publication of record for individual papers is online in the SPIE Digital Library.

SPIDigitalLibrary.org

Paper Numbering: Proceedings of SPIE follow an e-First publication model, with papers published first online and then in print and on CD-ROM. Papers are published as they are submitted and meet publication criteria. A unique, consistent, permanent citation identifier (CID) number is assigned to each article at the time of the first publication. Utilization of CIDs allows articles to be fully citable as soon as they are published online, and connects the same identifier to all online, print, and electronic versions of the publication. SPIE uses a six-digit CID article numbering system in which:

- The first four digits correspond to the SPIE volume number.
- The last two digits indicate publication order within the volume using a Base 36 numbering system employing both numerals and letters. These two-number sets start with 00, 01, 02, 03, 04, 05, 06, 07, 08, 09, 0A, 0B ... 0Z, followed by 10-1Z, 20-2Z, etc.

The CID number appears on each page of the manuscript. The complete citation is used on the first page, and an abbreviated version on subsequent pages. Numbers in the index correspond to the last two digits of the six-digit CID number.

Contents

xiii	Conference Committee
xv	Introduction
xvii	Remote sensing of snow hydrology: state of art and perspectives (Invited Presentation Summary) J. L. Foster NASA Goddard Space Flight Ctr. (United States)
xix	Exploiting the full potential of spectral observations for mapping vegetation biophysical variables: state of art and perspectives (Invited Presentation Summary) C. Atzberger, Univ. of Natural Resources and Life Sciences (Austria)
xxi	The evolution of airborne chemical and radiological remote sensing for emergency and natural disaster response (Plenary Summary) P. E. Lewis, National Geospatial-Intelligence Agency (United States)

SESSION 1 SNOW

8174 02	Remote sensing of snow cover and snow water equivalent for the historic February snowstorms in the Baltimore/Washington D.C. area during February 2010 (Invited Paper) [8174-01] J. L. Foster, D. Hall, NASA Goddard Space Flight Ctr. (United States); G. Riggs, NASA Goddard Space Flight Ctr. (United States) and SSAI (United States)
8174 03	Snow evolution in Sierra Nevada (Spain) from an energy balance model validated with Landsat TM data [8174-02] J. Herrero, Univ. de Granada (Spain); M. J. Polo, Univ. de Córdoba (Spain); M. A. Losada, Univ. de Granada (Spain)
8174 04	Characterization of snow pack over Pyrenees using remote sensed data for runoff modeling [8174-03] L. Moreno, A. Reppucci, X. Banque, Starlab (Spain); M. J. Escorihuela, IsardSAT (Spain); M. Aran, Applus Agroambiental, S.A. (Spain); R. Sanchez Diezma, HYDS (Spain)

SESSION 2 HYDROLOGY I

8174 05	Local effects on the water balance in flood plains induced by dam filling in Mediterranean environments [8174-04] C. Aguilar, M. J. Polo, Univ. de Córdoba (Spain)
8174 06	The use of LiDAR-derived high-resolution DSM and intensity data to support modelling of urban flooding [8174-05] M. Aktaruzzaman, T. G. Schmitt, Univ. Kaiserslautern (Germany)

8174 07 **Tracking, sensing and predicting flood wave propagation using nomadic satellite communication systems and hydrodynamic models** [8174-06]
R. Hostache, P. Matgen, L. Giustarini, C. Tailliez, J.-F. Iffly, Ctr. de Recherche Public - Gabriel Lippmann (Luxembourg)

8174 08 **River discharge estimation through MODIS data** [8174-07]
A. Tarpanelli, L. Brocca, Consiglio Nazionale delle Ricerche (Italy); T. Lacava, Consiglio Nazionale delle Ricerche (Italy) and Univ. degli Studi della Basilicata (Italy); M. Faruolo, F. Melone, T. Moramarco, Consiglio Nazionale delle Ricerche (Italy); N. Pergola, Consiglio Nazionale delle Ricerche (Italy) and Univ. degli Studi della Basilicata (Italy); V. Tramutoli, Univ. degli Studi della Basilicata (Italy) and Consiglio Nazionale delle Ricerche (Italy)

SESSION 3 HYDROLOGY II

8174 09 **Predicting soil erosion under land-cover area and climate changes using the revised universal soil loss equation** [8174-09]
S. Park, C. Jin, C. Choi, Pukyong National Univ. (Korea, Republic of)

8174 0A **What perspective in remote sensing of soil moisture for hydrological applications by coarse-resolution sensors** [8174-10]
L. Brocca, F. Melone, T. Moramarco, Consiglio Nazionale delle Ricerche (Italy); W. Wagner, Technische Univ. Wien (Austria)

8174 0B **Interception modeling with vegetation time series derived from Landsat TM data** [8174-12]
M. J. Polo, Univ. de Córdoba (Spain); A. Díaz-Gutiérrez, M. P. González-Dugo, Andalusian Institute of Agricultural and Fishery Formation (Spain)

8174 0C **Retrieving rainfall fields through tomographic processing applied to radiobase network signals** [8174-14]
F. Cuccoli, L. Facheris, S. Gori, Univ. degli Studi di Firenze (Italy); L. Baldini, Istituto di Scienze dell'Atmosfera e del Clima (Italy)

SESSION 4 CHANGE DETECTION

8174 0D **Damage estimation on agricultural crops by a flood** [8174-15]
N. del Carmen Silva-Aguila, A. López-Caloca, J. L. Silván-Cárdenas, Ctr. de Investigación en Geografía y Geomática (Mexico)

8174 0E **Evaluation of time-series and phenological indicators for land cover classification based on MODIS data** [8174-16]
F. Vuolo, Univ. of Natural Resources and Life Sciences (Austria); K. Richter, Ludwig-Maximilians-Univ. München (Germany); C. Atzberger, Univ. of Natural Resources and Life Sciences (Austria)

8174 0F **A multispectral multiplatform based change detection tool for vegetation disturbance on Irish peatlands** [8174-17]
J. O'Connell, J. Connolly, N. M. Holden, Univ. College Dublin (Ireland)

8174 0G **Mapping peatland disturbance in Ireland: an object oriented approach** [8174-18]
J. Connolly, N. M. Holden, Univ. College Dublin (Ireland)

SESSION 5 GENERAL APPLICATIONS I

- 8174 0H **Unmanned Aerial Vehicle (UAV) operated spectral camera system for forest and agriculture applications (Invited Paper)** [8174-19]
H. Saari, VTT Technical Research Ctr. of Finland (Finland); I. Pellikka, Univ. of Jyväskylä (Finland); L. Pesonen, MTT Agrifood Research Finland (Finland); S. Tuominen, Finnish Forest Research Institute (Finland); J. Heikkilä, Pieneering Ltd. (Finland); C. Holmlund, J. Mäkyneen, K. Ojala, T. Antila, VTT Technical Research Ctr. of Finland (Finland)
- 8174 0I **Automated object detection of climate tracers in remote-sensing data** [8174-20]
L. Tyrallová, Univ. Potsdam (Germany)
- 8174 0J **Passive unmanned sky spectroscopy for remote bird classification** [8174-21]
P. Lundin, M. Brydegaard, Lund Univ. (Sweden); L. Cocola, Lund Univ. (Sweden) and Univ. of Padova (Italy); A. Runemark, S. Åkesson, Lund Univ. (Sweden); S. Svanberg, Lund Univ. (Sweden), Zhejiang Univ. (China), and South China Normal Univ. (China)
- 8174 0K **Methods and potentials for using satellite image classification in school lessons** [8174-22]
K. Voß, R. Goetzke, H. Hodam, Rheinische Friedrich-Wilhelms-Univ. Bonn (Germany)

SESSION 6 IRRIGATION AND ENERGY BALANCE I

- 8174 0L **Improvement of the triangle method for soil moisture evaluation by adding a third index: albedo or cellulose absorption index** [8174-23]
J.-C. Krapez, ONERA (France); A. Olioso, Institut National de la Recherche Agronomique (France)
- 8174 0M **Energy balance modeling of agricultural areas by remote sensing** [8174-24]
A. Andreu, M. P. González-Dugo, IFAPA (Spain); M. J. Polo, Univ. de Córdoba (Spain); F. L. M. Padilla, P. Gavilán, IFAPA (Spain)
- 8174 0N **Relationships between high resolution RapidEye based fPAR and MODIS vegetation indices in a heterogeneous agricultural region** [8174-25]
S. Fritsch, Univ. of Würzburg (Germany) and Zentrum für Entwicklungsforschung (Germany); M. Machwitz, C. Conrad, Univ. of Würzburg (Germany); S. Dech, Deutsches Zentrum für Luft- und Raumfahrt e.V. (Germany)
- 8174 0O **Comparing actual evapotranspiration and plant water potential on a vineyard** [8174-26]
A. Maltese, C. Cammalleri, F. Capodici, G. Ciraolo, F. Colletti, G. La Loggia, T. Santangelo, Univ. degli Studi di Palermo (Italy)
- 8174 0P **Estimating evapotranspiration of riparian vegetation using high resolution multispectral, thermal infrared and lidar data** [8174-27]
C. M. U. Neale, H. Geli, S. Taghvaeian, A. Masih, R. T. Pack, Utah State Univ. (United States); R. D. Simms, M. Baker, J. A. Milliken, S. O'Meara, A. J. Witherall, U.S. Bureau of Reclamation (United States)

SESSION 7 VEGETATION

- 8174 0Q **Why confining to vegetation indices? Exploiting the potential of improved spectral observations using radiative transfer models (Invited Paper) [8174-28]**
C. Atzberger, Univ. of Natural Resources and Life Sciences (Austria); K. Richter, Ludwig-Maximilians-Univ. München (Germany); F. Vuolo, Univ. of Natural Resources and Life Sciences (Austria); R. Darvishzadeh, Shahid Beheshti Univ. (Iran, Islamic Republic of); M. Schlerf, International Institute for Geo-Information Science and Earth Observation (Netherlands)
- 8174 0R **Goodness-of-fit measures: what do they tell about vegetation variable retrieval performance from Earth observation data [8174-29]**
K. Richter, T. B. Hank, Ludwig-Maximilians-Univ. München (Germany); C. Atzberger, Univ. of Natural Resources and Life Sciences (Austria); W. Mauser, Ludwig-Maximilians-Univ. München (Germany)
- 8174 0S **Chlorophyll and soil effects on vegetation colorimetric characters [8174-31]**
R. Kancheva, G. Georgiev, Space and Solar-Terrestrial Research Institute (Bulgaria)
- 8174 0T **Multispectral vegetative canopy parameter retrieval [8174-32]**
C. C. Borel, D. J. Bunker, Air Force Institute of Technology (United States)
- 8174 0V **Evaluation of remotely sensed DMP product using multi-year field measurements of biomass in West Africa [8174-34]**
F. Nutini, Consiglio Nazionale delle Ricerche (Italy) and Univ. degli Studi di Milano (Italy); D. Stroppiana, M. Boschetti, P. A. Brivio, Consiglio Nazionale delle Ricerche (Italy); E. Bartholomé, European Commission Joint Research Ctr. (Italy); G. Beye, Ctr. de Suivi Ecologique (Senegal)

SESSION 8 AGRICULTURE I

- 8174 0X **Evaluation of red-edge spectral information for biotope mapping using RapidEye [8174-36]**
M. Bindel, S. Hese, C. Berger, C. Schmullius, Friedrich-Schiller-Univ. Jena (Germany)
- 8174 0Z **Remote mapping of susceptible areas to soil salinity, based on hyperspectral and geochemical data in the southern part of Tunisia [8174-38]**
M. Bouaziz, Technische Univ. Bergakademie Freiberg (Germany) and National School of Engineers of Sfax (Tunisia); R. Gloaguen, Technische Univ. Bergakademie Freiberg (Germany); B. Samir, National School of Engineers of Sfax (Tunisia)

SESSION 9 AGRICULTURE II

- 8174 11 **Investigating agro-drought in the Lower Mekong Basin using MODIS NDVI and land surface temperature data [8174-40]**
N. T. Son, C. F. Chen, C. R. Chen, National Central Univ. (Taiwan)

- 8174 12 **Potentials of RapidEye time series for improved classification of crop rotations in heterogeneous agricultural landscapes: experiences from irrigation systems in Central Asia** [8174-41]
C. Conrad, Julius-Maximilians-Univ. Würzburg (Germany); M. Machwitz, Zentrum für Entwicklungsforschung (Germany); G. Schorcht, F. Löw, Julius-Maximilians-Univ. Würzburg (Germany); S. Fritsch, Julius-Maximilians-Univ. Würzburg (Germany) and Zentrum für Entwicklungsforschung (Germany); S. Dech, Julius-Maximilians-Univ. Würzburg (Germany) and Deutsches Zentrum für Luft- und Raumfahrt e.V. (Germany)
- 8174 13 **Spectral and agronomical indicators of crop yield** [8174-42]
R. Kancheva, G. Georgiev, Space and Solar-Terrestrial Research Institute (Bulgaria)
- 8174 14 **Monitoring rice growing areas in the Lower Mekong subregion from MODIS satellite imagery** [8174-43]
C. R. Chen, C. F. Chen, N. T. Son, National Central Univ. (Taiwan)
- 8174 15 **Plastic covered vineyard extraction from airborne sensor data with an object-oriented approach** [8174-44]
E. Tarantino, A. Aiello, Politecnico di Bari (Italy)

SESSION 10 ESTUARINE, COASTAL, AND INLAND WATERS

- 8174 16 **Monitoring Mediterranean marine pollution using remote sensing and hydrodynamic modelling** [8174-45]
G. La Loggia, F. Capodici, G. Ciruolo, Univ. degli Studi di Palermo (Italy); A. Drago, Univ. of Malta (Malta); A. Maltese, Univ. degli Studi di Palermo (Italy)
- 8174 18 **Jellyfish prediction of occurrence from remote sensing data and a non-linear pattern recognition approach** [8174-47]
A. Albajes-Eizagirre, L. Romero, A. Soria-Frisch, Starlab Barcelona S.L. (Spain); Q. Vanhellefont, Institut Royal des Sciences Naturelles de Belgique (Belgium)
- 8174 19 **Characterizing the spectral signatures and optical properties of dams in Cyprus using field spectroradiometric measurements** [8174-48]
C. Papoutsas, D. G. Hadjimitsis, D. Alexakis, Cyprus Univ. of Technology (Cyprus)
- 8174 1A **Development of Japanese inland water surface temperature database using ASTER thermal infrared imagery** [8174-49]
H. Tonooka, M. Hirayama, Ibaraki Univ. (Japan)
- 8174 1B **Preliminary work of mangrove ecosystem carbon stock mapping in small island using remote sensing: above and below ground carbon stock mapping on medium resolution satellite image** [8174-50]
P. Wicaksono, P. Danoedoro, H. Hartono, Univ. Gadjah Mada (Indonesia); U. Nehren, L. Ribbe, Fachhochschule Köln (Germany)

POSTER SESSION: HYDROLOGY

- 8174 1C **A diachronic analysis of estuarine turbidity due to a flood following an extreme rainfall event** [8174-52]
A. Maltese, F. Capodici, G. Ciraolo, Univ. degli Studi di Palermo (Italy); A. Granata, Agenzia Regionale per la Protezione Ambientale (Italy); G. La Loggia, Univ. degli Studi di Palermo (Italy)

POSTER SESSION: AGRICULTURE

- 8174 1E **Use of imaging spectroscopy to assess different organic carbon fractions of agricultural soils** [8174-56]
M. Vohland, M. Harbich, O. Schmidt, Univ. Trier (Germany); T. Jarmer, Univ. Osnabrueck (Germany); C. Emmerling, S. Thiele-Bruhn, Univ. Trier (Germany)

POSTER SESSION: VEGETATION

- 8174 1F **An improvement of satellite-based algorithm for gross primary production estimation optimized over Korea** [8174-58]
K.-J. Pi, K.-S. Han, I.-H. Kim, S.-I. Kim, M.-J. Lee, Pukyong National Univ. (Korea, Republic of)
- 8174 1G **Assessment of tundra-taiga boundary changes using MODIS LAI data** [8174-59]
M. Lee, K.-S. Han, K.-J. Pi, S.-I. Kim, I.-H. Kim, Pukyong National Univ. (Korea, Republic of)
- 8174 1H **On the influences of vegetation biomass on COSMO-Skymed X-band** [8174-60]
F. Capodici, Univ. degli Studi di Napoli Federico II (Italy) and Univ. degli Studi di Palermo (Italy); G. Ciraolo, Univ. degli Studi di Palermo (Italy); G. D'Urso, Univ. degli Studi di Napoli Federico II (Italy); G. La Loggia, A. Maltese, Univ. degli Studi di Palermo (Italy)
- 8174 1I **Airborne spray drift measurement using passive collectors and lidar systems** [8174-61]
E. Gregorio, Univ. de Lleida (Spain); F. Solanelles, Generalitat de Catalunya (Spain); F. Rocadenbosch, Univ. Politècnica de Catalunya (Spain); J. R. Rosell, R. Sanz, Univ. de Lleida (Spain)

POSTER SESSION: THERMAL IR

- 8174 1J **Spatial analysis of LST in relation to surface moisture and NDBI using landsat imagery in Cheongju city** [8174-62]
S.-I. Na, J.-H. Park, S.-C. Baek, Chungbuk National Univ. (Korea, Republic of)
- 8174 1M **The *Rhynchophorus ferrugineus* disease of *Phoenix canariensis*: early detection through proximity thermal sensing** [8174-65]
C. Cammalleri, Univ. degli Studi di Palermo (Italy); F. Capodici, Univ. degli Studi di Napoli Federico II (Italy); G. Ciraolo, G. Filardo, G. La Loggia, A. Maltese, Univ. degli Studi di Palermo (Italy)

POSTER SESSION: ESTUARINE, COASTAL AND INLAND WATERS

- 8174 1N **Spatio-temporal variability analysis of the Douro River plume through MERIS data for one hydrological year** [8174-67]
A. Teodoro, H. Almeida, Univ. do Porto (Portugal)
- 8174 1S **Multi-temporal spatial pattern analysis of Tulum's tropical coastal landscape** [8174-72]
S. C. Ramírez-Forero, A. López-Caloca, J. L. Silván-Cárdenas, Ctr. de Investigación en Geografía y Geomática (Mexico)

POSTER SESSION: CHANGE DETECTION

- 8174 1T **A statistical model for the selection of ground observations of solar radiation: an application in producing a five-year dataset of radiation maps on Italian territory through correction of MSG-derived data** [8174-74]
L. Campo, F. Castelli, Univ. degli Studi di Firenze (Italy)
- 8174 1U **An ecosystem service value assessment of land-use change in South Korea using remote sensing data and geographic information system** [8174-75]
Y. Jung, H. Ahn, C. Choi, Pukyong National Univ. (Korea, Republic of)

POSTER SESSION: IRRIGATION AND ENERGY BALANCE

- 8174 1W **Scaling from instantaneous remote-sensing-based latent heat flux to daytime integrated value with the help of SiB2** [8174-77]
Y. Song, Cold and Arid Regions Environmental and Engineering Research Institute (China) and Institute of Earth Environment (China); M. Ma, X. Li, Cold and Arid Regions Environmental and Engineering Research Institute (China); X. Wang, Cold and Arid Regions Environmental and Engineering Research Institute (China) and Graduate Univ. of Chinese Academy of Sciences (China)

POSTER SESSION: GENERAL APPLICATIONS

- 8174 1X **Spatial distribution of carbon dioxide absorption and emission in Chungcheongbuk-do, South Korea using RS and GIS method** [8174-78]
J.-K. Park, S. Na, J.-H. Park, Chungbuk National Univ. (Korea, Republic of)
- 8174 1Y **An integration system for the collocation of polar and geostationary satellite observations** [8174-79]
H. Sun, Perot Systems Government Service (United States); W. Wolf, National Oceanic and Atmospheric Administration (United States); T. King, Perot Systems Government Service (United States); C. Barnet, M. Goldberg, National Oceanic and Atmospheric Administration (United States)

POSTER SESSION: RS FOR AGRICULTURE, ECOSYSTEMS, AND HYDROLOGY

- 8174 20 **A remote sensing technique for the assessment of stable interannual dynamical patterns of vegetation** [8174-82]
M. Y. Chernetskiy, A. P. Shevyrnogov, Institute of Biophysics (Russian Federation) and Siberian Federal Univ. (Russian Federation); N. F. Ovchinnikova, V.N. Sukachev Institute of Forest of SB RAS (United States)
- 8174 22 **GPP estimation over Heihe River Basin, China** [8174-84]
X. Wang, Cold and Arid Regions Environmental and Engineering Research Institute (China) and Graduate Univ. of Chinese Academy of Sciences (China); M. Ma, X. Li, X. Han, Y. Ran, G. Huang, Cold and Arid Regions Environmental and Engineering Research Institute (China); Y. Song, Cold and Arid Regions Environmental and Engineering Research Institute (China) and Graduate Univ. of Chinese Academy of Sciences (China); J. Tan, Cold and Arid Regions Environmental and Engineering Research Institute (China)
- 8174 23 **Validation of MODIS land surface temperature products using ground measurements in the Heihe River Basin, China** [8174-85]
W. Yu, Cold and Arid Regions Environmental and Engineering Research Institute (China) and Graduate Univ. of Chinese Academy of Sciences (China); M. Ma, Cold and Arid Regions Environmental and Engineering Research Institute (China); X. Wang, Y. Song, Cold and Arid Regions Environmental and Engineering Research Institute (China) and Graduate Univ. of Chinese Academy of Sciences (China); J. Tan, Cold and Arid Regions Environmental and Engineering Research Institute (China)
- 8174 24 **Agricultural land cover classification using rapideye satellite imagery in South Korea - first result -** [8174-88]
H. O. Kim, J. M. Yeom, Y. S. Kim, Korea Aerospace Research Institute (Korea, Republic of)
- 8174 26 **A computer simulation model to compute the radiation transfer of mountainous regions** [8174-91]
Y. Li, F. Zhao, R. Song, Beijing Univ. of Aeronautics and Astronautics (China)
- 8174 27 **Mutual influence between climate and vegetation cover through satellite data in Egypt** [8174-92]
M. A. El-Shirbeny, M. A. Aboelghar, S. M. Arafat, National Authority for Remote Sensing and Space Sciences (Egypt); A.-G. M. El-Gindy, Ain Shams Univ. (Egypt)
- 8174 28 **DEM densification using SFS with single multi-spectral satellite image** [8174-94]
Z. Chen, T. Sun, Q. Qin, Wuhan Univ. (China); H. Zhang, The Second Institute of Oceanography (China)
- 8174 29 **Periodicity analysis of NDVI time series and its relationship with climatic factors in the Heihe River Basin in China** [8174-95]
H. B. Han, Anhui Agricultural Univ. (China); M. G. Ma, Cold and Arid Regions Environmental and Engineering Research Institute (China); P. Yan, Anhui Agricultural Univ. (China); Y. Song, Cold and Arid Regions Environmental and Engineering Research Institute (China)

- 8174 2C **Angular and polarization measurements of snow and bare soil microwave reflective and emissive characteristics by ka-band (37GHz), combined scatterometer-radiometer system** [8174-102]
M. L. Grigoryan, A. K. Arakelyan, A. K. Hambaryan, A. A. Arakelyan, ECOSERV Remote Observation Ctr. Co. Ltd. (Armenia)
- 8174 2E **Estimation of water budget by remote sensing in Taihu Basin, China** [8174-104]
X. Zhao, Y. Liu, Nanjing Institute of Geography and Limnology (China)
- 8174 2G **A radiosity-based model to compute the radiation transfer of soil surface** [8174-106]
F. Zhao, Y. Li, Beijing Univ. of Aeronautics and Astronautics (China)
- 8174 2H **Validation of collection 5 MODIS LAI product by scaling-up method using field measurements** [8174-107]
H. Xue, Beijing Normal Univ. (China) and Henan Polytechnic Univ. (China); J. Wang, C. Ping, Beijing Normal Univ. (China)

Author Index

Conference Committee

Symposium Chair

Karin Stein, Fraunhofer-Institut für Optronik, Systemtechnik und Bildauswertung (Germany)

Symposium Cochair

Charles R. Bostater, Florida Institute of Technology (United States)

Conference Chairs

Christopher M. U. Neale, Utah State University (United States)

Antonino Maltese, Università degli Studi di Palermo (Italy)

Conference Cochair

Katja Richter, Ludwig-Maximilians-Universität München (Germany)

Program Committee

Guido D'Urso, Università degli Studi di Napoli Federico II (Italy)

Goffredo La Loggia, Università degli Studi di Palermo (Italy)

Francesco Vuolo, University of Southampton (United Kingdom)

Session Chairs

- 1 Snow
Christopher M. U. Neale, Utah State University (United States)
- 2 Hydrology I
Goffredo La Loggia, Università degli Studi di Palermo (Italy)
- 3 Hydrology II
Christopher M. U. Neale, Utah State University (United States)
- 4 Change Detection
Katja Richter, Ludwig-Maximilians-Universität München (Germany)
- 5 General Applications I
Antonino Maltese, Università degli Studi di Palermo (Italy)
- 6 Irrigation and Energy Balance I
Antonino Maltese, Università degli Studi di Palermo (Italy)

- 7 Vegetation
Francesco Vuolo, Universität für Bodenkultur Wien (Austria)
- 9 Agriculture I
Antonino Maltese, Università degli Studi di Palermo (Italy)
- 10 Agriculture II
Katja Richter, Ludwig-Maximilians-Universität München (Germany)
- 11 Estuarine, Coastal, and Inland Waters
Christopher M. U. Neale, Utah State University (United States)

Introduction

This proceedings volume contains papers presented during the conference on Remote Sensing for Agriculture, Ecosystems, and Hydrology XIII. The conference was part of the 18th International Symposium on Remote Sensing sponsored by SPIE—The International Society for Optical Engineering. The symposium was held at the Clarion Congress Hotel Prague, Prague, Czech Republic, from 19th to 21th of September 2011.

The conference is dedicated to providing rapid dissemination of scientific and technical information, and attracted scientists and professionals from throughout Europe, Africa, Asia, and the Americas. Approximately 45 oral and 30 poster presentations were given, covering a broad range of topics in the field of remote sensing applications in environmental science.

The program was organized according to major themes, with 10 sessions on Agriculture: Irrigation and Energy Balance and Agriculture (2); Ecosystems: Estuarine, Coastal and Inland Waters; Vegetation and Change detection; Hydrology: Snow and Hydrology (2). The poster presentations also had good representation from the abovementioned themes. The presentations described both fundamental and applications-based research activities from modelling, to laboratory and field experiments, to operational applications. The oral program also included three invited presentations: James L. Foster of NASA Goddard Space Flight Ctr. (USA) gave a presentation on the subject *"Remote sensing of snow cover and snow water equivalent for the historic snowstorms in the Baltimore/Washington area during February 2010"* within a Snow session; Heikki K. Saari of VTT Technical Research Ctr. of Finland (Finland) gave a presentation on the subject *"Unmanned aerial vehicle (UAV) operated spectral camera system for forest and agriculture applications"* within General Application session; Clement Atzberger of University of Natural Resources and Life Sciences (BOKU) Wien, (Austria) gave a presentation on the subject *"Why confining to vegetation indices? Exploiting the potential of improved spectral observations"* within a Vegetation session; Short reports pointing out the state of art and perspectives in the research fields of the invited talks are reported below.

We extend our thanks to the co-editor Katja Richter of Ludwig-Maximilians-Univ. München, and Session chairs Francesco Vuolo of Univ. of Univ. für Bodenkultur Wien (Austria) and Goffredo La Loggia of Univ. degli Studi di Palermo, and to the presenters for their efforts and to the participants for their insightful questions and discussions. Special thanks are also due to the host city for the excellent venue and to all the SPIE organizational staff for their support prior to, during, and after the symposium. We look forward to an even more successful conference in 2012.

Christopher M. U. Neale
Antonino Malfese
Katja Richter

Remote sensing of snow hydrology: state of art and perspectives

Snow plays an important role in the global energy and water budgets, as a result of its high albedo and thermal and water storage properties. Snow is also the largest varying landscape feature of the Earth's surface. For example, in North America, the snow cover extent may vary from greater than 50% to less than 5% in the course of six months, and the snow water equivalent (SWE) of mid-latitude snowpacks can be reduced by as much as 100 mm in less than six days. Furthermore, snow depth and SWE, as well as snow cover extent, are important contributors to both local and remote climate. Thus, knowledge of snow extent and SWE are important for climate change studies and applications such as flood forecasting.

Melting snow contributes much of the water required for drinking, agriculture and, industry in many regions of the world. In the western U.S., upwards of 70% of the water supply is derived from snowmelt. In India, Pakistan, Afghanistan, and Nepal snow and ice melt from the Hindu Kush and Himalayan ranges are a vital resource for nearly 1 billion people. The ability to characterize snow storage more accurately at the drainage basin scale is crucial for improved management of our precious water resources. Snowmelt data are needed in hydrological models to improve flood control and irrigation. Also, SWE, snow cover, and melt onset are critically needed parameters for climate modeling and the initialization of forecasts at weather and seasonal time scales.

There is a heritage of 50 years of remotely sensed snow observations utilizing the visible wavelengths and 35 years utilizing passive microwave observations. We now have sufficient years of observation to develop robust datasets so that meaningful trends in snow cover extent can be detected and related to climate fluctuations. Additionally, we now have a much better understanding of sensor retrieval errors. This is crucial in order to correctly interpret and evaluate the observations and to successfully assimilate them into numerical models.

Despite its importance, successfully forecasting and modeling snowmelt is a challenge since the knowledge of snow physics is imperfect, even the most sophisticated models are simplifications and because errors still exist in the model forcing data. Moreover, the natural spatial and temporal variability of snow cover is characterized at space and time scales below those typically represented by models. Snow model initialization based on model spin-up will be affected by these errors. As we become more adept and comfortable in assimilating snow observation products into Land Surface Models (LSMs), the effects of model initialization error will likely be reduced with the end result being models that more accurately and reliably resemble observations.

James L. Foster
Invited Speaker
NASA-GSFC
Greenbelt, United
States of America

Exploiting the full potential of spectral observations for mapping vegetation biophysical variables: state of art and perspectives

Continuous fields of vegetation biophysical variables (e.g. LAI) are required in numerous applications (e.g. precision agriculture, irrigation management, SVAT models). Proposed methods for mapping biophysical variables range from empirical techniques mainly based on vegetation indices (VI) to physically based techniques employing radiative transfer models. Albeit numerous experiments found close relationships between VI and important biophysical variables, well known shortcomings and drawbacks remain:

Under-determined inverse problem: Even when looking only at relatively uniform agricultural crops, the measured canopy spectral reflectance (visible to SWIR) is influenced by at least by 5-10 structural and optical properties for a given illumination and observation geometry. The main structural properties affecting the canopy signature are LAI, leaf orientation (ALA) and leaf clumping. The leaf optical properties are mainly driven by the leaf chlorophyll and water content, and the leaf structure. In addition, soil optical properties are known to change with surface wetness and roughness. For this reason, attempts to define a close relation between a (2-band) VI and for example LAI, can only work well under specific conditions, where all other parameters are more or less constant. Generally, 2-band vegetation indices such as NDVI only minimize soil background influences.

Sub-optimum use of available spectral information: Most sensors record spectral data in more than two bands. Hence, the available spectral information of multi-spectral sensors and imaging spectrometers is not fully exploited when vegetation indices are used for mapping vegetation biophysical variables. Important information is ignored when using only a few (e.g. 2) wavebands for modeling LAI or other biophysical variables. If empirical techniques are to be used, it is certainly more appropriate to use full spectrum methods such as Partial Least Square (PLS) regression. Full spectrum methods are at the same time less sensitive to (Gaussian) noise as compared to VI.

Model over-specialisation: Vegetation indices can be defined and fine tuned to different environmental settings, vegetation types, and measurement conditions. This advantage, however, often leads to over-specialized models which are only valid for the specific training data base used for model calibration. For a given training data base it can be shown that the 'optimum' functional form of the VI and selected 'optimum' wavelengths are very sensitive to measurement uncertainties and calibration strategies (e.g. cross-validation, bootstrapping). Therefore, developed models can hardly be reproduced. Internal validation strategies generally yield over-optimistic performance statistics.

Lack of transferability: In most spectral regions, even relatively small spectral shifts lead to changes in the observed canopy spectral reflectance. At the same time,

vegetation canopies are strong anisotropic scatterers, where the recorded signal depends on the actual sun/view geometry. Hence, empirical models developed for a given measurement geometry and sensor (e.g. with specific band widths and spectral response functions) can generally not be transferred to other sensors and measurement conditions. In practical terms this implies high sampling costs (man power) for large area mappings.

Most of the above-presented drawbacks cannot be overcome whatever mathematical transformation is applied to the data. Hence, constantly proposing new vegetation indices (e.g. functional forms and waveband combinations) is probably unproductive. Instead, research should focus on full spectrum methods and in particular physically based approaches, which provide an explicit connection between the recorded signal and the sought biophysical variables. In this way, sensor characteristics and measurement conditions are implicitly accounted for. Methods built on physical principles also better generalize across different environmental settings and are less prone to over-specialisation. The ill-posed inverse problem can be (partly) overcome by imposing constraints on the inverse problem. For example, several studies demonstrated the benefits of exploiting spatial variations in the signal (e.g. object-based inversions or patch-ensemble approaches). In addition, temporal constraints can be used to further regularize the inverse problem and to fully exploit the intrinsic dimensionality of the recorded signal.

Clement Atzberger
Invited Speaker
University of Natural
Resources and Life
Sciences (BOKU)
Wien, Austria

Plenary Summary

The Evolution of Airborne Chemical and Radiological Remote Sensing For Emergency and Natural Disaster Response

Summary of the September 19, 2011 SPIE Remote Sensing Plenary Session Presentation by

Paul E. Lewis

National Geospatial-Intelligence Agency, United States of America

First responders, joint operations centers, and recovery and remediation personnel consider timely and affordable airborne chemical, radiological, imagery analysis, and related mapping products essential in the formulation of a complete understanding of an incident and its potential impact on adjacent communities, and for recovery and remediation. Airborne remote sensing provides the flexibility to produce incident specific products and conduct over-flights at the frequencies needed to provide timely and relevant information for recovery and remediation operations, optimization of resources during an event, and for the safety of emergency response personnel.

The utility of airborne chemical remote sensing became apparent to the EPA during a chemical plant explosion, which occurred in Sioux City, Iowa in December of 1994. The facility produced ammonium nitrate fertilizer, and also produced its own ammonia for use in the process. In late December an explosion occurred rupturing the main storage tank and spilling three million gallons of ammonia. This resulted in lethal vapor levels in and around the plant and created a plume of ammonia vapors estimated to be 35 miles long. Approximately 3,500 people were evacuated over a 50 square mile area. The EPA sent in vehicles with ground sampling crews dressed in Level A hazmat suits with 30 minute air packs to monitor the site. Due to heavy snow coverage on the ground and saturated soil conditions underneath the snow, all of the EPA vehicles became stuck. Ground sampling crews had to be rescued before air supplies ran out. Consequently, no monitoring of vapor levels was accomplished.

The lessons learned from responding to the chemical explosion in Sioux City, Iowa in 1994 prompted the EPA to begin evaluating the application of airborne remote sensing infrared and gamma ray spectroscopy for emergency responses involving chemical and radiological incidents. Concurrently, with the evaluation process to determine the performance and feasibility of implementing infrared and gamma ray spectroscopy in an airborne platform came the evolution of a set of core requirements for an airborne operational capability: Standoff chemical and gamma ray detection and identification with low false alarm rates; High resolution ortho-rectified day-night imagery; Airborne data collection under cloud ceilings; Rapid dispatch-wheels up in under one hour after activation; Automated data processing –real or near-real-time chemical data analysis; Direct integration of data and information to local incident commanders-local and federal joint operations centers; Data telemetry to and from the aircraft.

According to the EPA, in the United States there are approximately 123 facilities where a release of chemicals could threaten more than one million people. There are approximately 750 additional facilities where a chemical release could threaten more than a hundred thousand people.

In 2001, the EPA implemented the United States only civilian operational airborne chemical detection and identification capability called the Airborne Spectral Photometric Environmental Collection Technology (ASPECT) Program. Subsequently in 2003, the EPA and NGA agreed to collaborate in a cooperative research and development program focused on evolving the capabilities of the ASPECT Program to produce near-real-time state of the art chemical, radiological and imagery mapping emergency response products.

Plenary Summary

Airborne Spectral Photometric Environmental Collection Technology (ASPECT) Program
The United States Only Airborne
24/7 Operational CIVIL Emergency Response Chemical, Radiological, & Imaging Mapping Capability

The ASPECT model of operation combines an airborne operational remote sensing suite with a research and development support team to insure that analysis and products are validated and verified scientifically and are reviewed and checked before release. The research and development support team collaboration between the EPA and NGA to evolve the capabilities of the ASPECT Program has resulted in the following significant accomplishments: Near-real-time automated onboard chemical detection and identification of 78 chemical compounds with low false alarm rates; Near-real-time information on plume direction and concentrations; Automated software producing day/night ortho-rectified imagery rapid response maps; Automated software producing gamma ray survey information maps onboard the aircraft; Data and information telemetry to and from the aircraft facilitating turn-around times and seamless integration of vital situational awareness information from the aircraft to first responders or joint operation centers in 5 to 15 minutes.

Since 2001 the ASPECT Program has provided essential information during 115 emergency, disaster, and homeland security related incidents ranging from chemical plant explosions and train derailments to fires, floods, hurricanes, and special events. The ASPECT Program played key roles in providing essential information to first responders and joint operations centers in response to the following historical events: The Shuttle Columbia break up during re-entry over Texas in February of 2003; Hurricane Katrina in August of 2005; The Deepwater Horizon Oil Spill disaster in the Gulf of Mexico from April-August 2010.

Over the past decade in over 115 responses, the ASPECT program has demonstrated the utility of having timely, cost-effective operational airborne chemical and radiological remote sensing information integrated seamlessly into the local, state and federal emergency response and disaster recovery and remediation communities. What is needed next is the implementation of multiple aircraft strategically located throughout the United States so that ASPECT capabilities can be on the scene of a disaster or event in less than three hours.