NDEX

Α

Abbe sine condition, 162 Abbe V-value, 93 aberration, 74, 164, 166, 168, 174, 178, 181, 188, 191, 195, 211-213, 219-222, 227, 241, 245 angular ray, 221–224, 228, 247 of the Galilean telescope, 225 of infrared afocal system, 249 balancing, 82 bending for minimum spherical, 149, 158 chromatic, 204 coma, 76 higher-order, 82, 85, 169, 174, 177 201-202 longitudinal, 45 minimum spherical, 149-154, 158 monochromatic, 74 of centered optical systems, 70 off axis, 161, 165, 257, 260 Seidel, 107 pupil, 81 ray, 65 secondary, 82 Seidel, 44, 74 monochromatic, 233 spherical, 74-76, 82, 203, 234-241, 249,253-259, 262, 265-268, 272, 276 third order, 27, 74-75, 82-86, 102, 107, 113, 202 transverse ray, 45-46, 66-68, 83-85, 131, 134, 139-159 wavefront, 45, 55–57, 63–65, 68 achromatic doublet, 91-97, 167, 183

achromatized Ramsden, 214 Schmidt camera, 268 afocal system, 29, 81, 164, 209-212, 219, 224, 227, 247–249 angle of incidence, 2-3, 39, 42, 48, 53, 105–109, 254 angle of refraction, 2, 48, 53 angular magnification, 28-30, 81, 210, 219, 226 aperture stop, 6–8, 26, 41, 45, 97, 115-16, 142, 153, 159-161, 165, 209, 212, 267, 270-272 aplanatic surface, 122-125, 149, 164, 170, 195 aspheric infrared germanium lens, 234-236 Petzval lens, 241 plate, 159–161, 267 surface, 57, 120, 128, 157–162, 233-236, 239, 241-242, 253, 256, 258, 259, 261, 262, 263, 264, 266, 268, 269, 271 refracting, 164, 254 astigmatism, 72-74, 77-79, 85, 107, 112, 119, 131–135, 142, 152-154, 158-159, 165, 167-171, 179, 183-188, 194, 203, 212–221, 234, 240, 249, 255-257, 260-263, 267, 270, 276 asymmetric, 205 axial color, 92, 97, 111–114, 145, 151-152, 157-159, 165, 167, 169-171, 179, 214-215, 233-234, 241, 259, 276 axicon, 14

В

back focal length, 24–25, 200 baffles, 255 beam expanders, 28, 226 balancing aberration, 82 astigmatism, 85 coma, 83 distortion, 86 bending for minimum spherical aberration, 148, 156

С

Canzek-Mangin system, 276–278 Cassegrain telescope, 163 field corrector for, 257 cardinal points, 18, 32 catadioptric systems, 130, 253-255 cemented doublet, 151, 170-171, 177-178, 182, 183, 187-189, 214, 268 chief ray, 7, 26–27, 40–42, 44, 45, 55-56, 63, 67, 78-87, 97, 101-102, 107-108, 112-115, 126, 135–136, 139, 146, 157, 201, 212–211, 217, 221–222 chromatic aberration, 204 coma, 72-73, 74-76, 86, 88, 142, 149, 152-153, 158, 162-166, 170, 184-186, 194-195, 199-205, 213, 216, 255–262, 267, 270, 276 aberration, 76 corrector for tranverse, 260 balancing, 83 corrector for a parabolic mirror, 259 tangential, 241 concentric, 149, 165 conic constant, 11, 14 conjugate factor, 140-142, 150 finite, 65, 79 infinite, 65, 80 parameter, 152 points, 20, 21 shift effects, 136–137

Conrady chromatic aberration formula, 97–100, 111 convergence angle, 6 Cooke triplet, 114, 199–204 coordinate system, 1, 50, 61 crown glass, 151, 170–172 crown-first doublet, 171–172, 175 curvature, 9, 14, 23, 39, 58, 102, 110–112, 120–121, 126–128, 139, 155, 165, 200, 254–255, 267, 270–272 curved field, 186, 192, 217

D

defocus, 69, 77, 81, 83-84, 221, 227 diffraction, 64, 68, 75, 87 limit, 90, 101 direction cosines, 50-53, 60-61 dispersion, 91–97, 122, 145, 157, 171, 204–205, 268 low, 232–233 crown, 233 partial, 95-96 distortion, 79-82, 86, 108, 120, 127–131, 144, 159, 165, 195, 199–201, 204, 208, 212–213, 267 at finite conjugates, 80 fifth-order, 86 pupil, 135 third-order, 79

Ε

eccentricity, 41 ellipsoid, 10 hyperboloid, 13 entrance pupil, 7, 29–30, 65–71, 81, 113, 126, 135, 163–164, 210–211, 247, 249 equivalent focal length, 32 Erfle eyepiece, 217–218 exit pupil, 7–8, 29–30, 64, 71, 135, 164, 209–212, 221, 226, 247 eyepieces, 8, 209–217 eye relief, 209–211, 214, 217–219, 222, 226

F

f/numbers, 262 Fermat's principle, 4–5, 63, 99 field angle, 78-79, 82, 85-84, 88-89, 113, 129, 134, 165, 183–186, 191, 204–208, 210, 212–217, 221, 241, 249, 263, 272 corrector, 265, 276 for a Cassegrain telescope, 257 for a hyperbolic mirror, 265 for a parabolic mirror, 260 for a paraboloidal mirror, 260 for a Ritchey-Chrétien telescope, 263 - 264curvature, 72–74, 79–81, 88, 107, 110, 121–122, 127, 143, 153, 159, 162, 165, 168–169, 173, 178, 182, 184–185, 188–193, 201–204, 212, 247 217-221, 226, 254-255, 267, 270–272, 276 flattener, 128, 191–193 lens, 247, 267 stop, 8, 220, 224 fifth-order distortion, 86 finite conjugates, 29–30, 162 first-order optics, 74 paraxial, 92 first principal focus, 19-20, 30 plane, 19 point, 19 flint glass, 151, 170–171, 233 focal length, 24, 93–94, 97, 163, 182, 185, 200–202, 210–211, 214, 219, 222, 226, 253, 259 equivalent, 32 planes, 174, 207 sagittal, lines, 79

shift, 97, 146 tangential, lines, 79 focus, 19–20, 30, 92, 97, 128, 191 250, 257–268, 279 paraxial, 76, 80–81 f-theta lenses, 81

G

Galilean telescope, 224, 225, 226 Gauss doublet, 177, 179-181 Gaussian optics, 17 region, 17 geometrical approximation, 88 modulation transfer function (MTF). 89 optics, 1, 64, 88, 101 wavefront, 55 germanium, 150, 153, 232–240, 244, 247Petzval lens, 240–242 silicon triplet, 245-246 triplet, 242-244 glass choice, 171, 177

Η

higher-order aberrations, 75, 82, 85, 169, 174, 177, 186, 201–202 astigmatism, 85 sagittal oblique spherical aberration (SOBSA), 188 tangential oblique spherical aberration (TOBSA), 188 hyperbolic mirror field corrector for, 265 hyperboloid, 10–14, 262, 265 hyperboloidal secondary mirror, 163, 255

I

image curvature, 186, 212 space, 18–22, 29–33, 55, 65, 81, 98, 116, 132, 141, 212 surface, 79, 207 infrared afocal system, 247–248 detectors, 247 materials, 249–250 wavelengths, 153 interstitium, 37

Κ

Kellner eyepiece, 214–215

L

Lagrange invariant, 25–32, 40–41, 107, 113, 141, 161, 169, 172, 175, 178, 180, 185, 201, 213–218, 221, 223, 225, 228, 235, 237, 239, 241, 243, 245, 248, 256, 258, 259, 261, 263, 264, 266, 268, 269, 271, 273, 275, 277, 280 lateral color, 97, 100, 111–112, 120–122, 129, 146, 153, 158–159, 165–166, 170, 195, 201, 211–217, 226, 260, 270 image shift, 74 longitudinal aberrations, 45 low dispersion, 232–233 crown, 233

Μ

magnification, 20, 27–30, 129, 137, 141, 148–199, 153, 156, 162–166 magnifiers, 226–227 Malus, 63 and Dupin theorem, 5–6 Maksutov-Bouwers, 272–274 Cassegrain system, 272 Mangin mirror, 255, 274, 276 system, 279 marginal ray, 7, 25–28, 39–42, 64, 102, 107–109, 114–117, 126, 135, 163, 184, 191, 221 meridian plane, 7, 67, 71–73, 78 rays, 188 microscope objective, 125–126, 149 minimum spherical aberration, 148–153, 156 bending for, 149, 158 mirror, 4, 71, 82, 253–262, 265–267, 270–276, 279–282 modulation transfer function (MTF), 86, 188 monochromatic aberrations, 74, 147 Seidel, 233

Ν

Narcissus effect, 249 Newton's conjugate distance equation, 30–31 nodal points, 18–21, 33, 37 normal to the surface, 2, 60 numerical aperture, 65, 68, 87, 90, 116, 126, 168–169, 219

0

object space, 7, 18-21, 29-33, 46, 65, 81, 98, 141, 233, 247-249 surface, 65 objective, 81, 85-86, 113, 168, 171, 183–185, 191, 194, 195, 200-202, 209-212, 219, 220–226, 247 off-axis aberrations, 107, 161, 165, 168, 257, 260 optical glass, 93–94 optical path difference (OPD), 55, 69, 98, 110 optical path length, 4–5, 22, 99 optical transfer function, 67, 86–87, 90 optics first-order, 74 Gaussian, 17

geometrical, 1, 64, 88, 101 paraxial, 17, 27, 45 overcorrected aberrations, 128

Ρ

parabolic mirror, 265 paraboloid, 10, 259, 267, 281-282 paraboloidal primary mirror, 163 field corrector for, 260 paraxial, chief ray, 26-27, 139, 161 first order, 92 focus, 76, 80-83 foci, 269 image plane, 70, 79, 85, 134 size. 83 marginal ray, 26, 139, 161 optics, 17, 27, 45 surface, 70, 86 ray, 21–25, 102, 105–106, 111, 116, 121–122, 139, 162, 164, 175 tracing, 23-26, 34 region, 45, 63 value, 188 partial dispersion, 95-96 Petzval lens, 158, 183-193, 205, 226 objective, 191 sum, 127, 130–134, 154–156, 159, 169, 184, 189–191, 194, 199–200, 249 surface, 132–135, 154 primary aberrations, 81 mirror, 163 principal planes, 19-20, 35-37 points, 18, 29, 33-35 pupil, 6-7, 91-93, 186-188, 206, 247, 267 aberrations, 81, 135–136 astigmatism, 135 coma, 135

distortion, 135 entrance, 29, 81 exit, 29 field curvature, 135 sheared, 88

Q

quadrics of revolution, 10

R

ray, 17–44, 97–100, 144, 150, 173, 178, 188, 192 abberated, 63, 66, angular, 221–224, 228 chief, 27, 40–41, 46, 55–56, 63, 67, 78-81, 97 coordinates, 64 marginal, 27, 40-41, 64 meridian, 75 parallel, 34, 39 paraxial, 17, 21, 23–26, 34, 38, 102, 105–107, 112, 118, 123 - 124chief, 26–27, 40–42, 44, 46, 55-57, 63, 67-68, 81-84, 97, 101–102, 108, 113, 116–117, 128, 138–139, 140, 147, 159 first order, 92 marginal, 26–28, 39, 42, 221 sagittal, 75 skew, 46-48, 55, 67 tracing, 45-61, 66, 97, 99, 101, 110, 116, 131, 160, 169, 255 at reflecting surfaces, 61 paraxial, 25–26 trigonometric, 47 transverse, 63–65 reference sphere, 55–56, 63–64, 102 - 104reflecting surface, 3, 18, 130, 254 refracting surface, 5, 21-23, 48, 57, 102, 162, 255 refraction invariant, 39-40, 114

refractive index, 1–6, 55, 91–100, 109, 125, 129, 139–141, 144, 149–155, 162, 171, 189, 194, 205, 232–233, 250, 267 relative pupil coordinates, 64 relative ray coordinates, 68 Ritchey-Crétien telescopes, 163, 262, 265 field corrector for, 263–264

S

sagittal astigmatism, 85 curve, 173, 188, 203 focal lines, 79 foci, 184 focus, 78 image surface, 133–134, 184, 188.207 plane, 134 rays, 75 section, 67-68, 75-78, 185 surface, 154 Schmidt cameras, 162, 267–270 field-flattened, 270 Schott optical glass, 94, 96, 151, 202 secondary aberration, 82 mirror, 163 spectrum, 94, 173, 182, 260 second principal, 19 Seidel aberrations, 44, 74, 101–138 140-142, 146-147, 152, 160, 161–162, 186, 194, 199–201, 205, 255, 275 monochromatic, 233 off-axis, 107 third-order, 162 analysis, 184 chromatic aberration coefficients, 162 coefficient, 185 for astigmatism, 142 for field curvature, 143 of axial color, 145

of coma. 142 of distortion, 144 of lateral color, 146 of spherical aberration, 142 difference formulae, 161 theory, 130, 139, 174, 199, 253 - 254separated lenses, 154 shape factor, 139–142, 147–148, 151-153, 200-201 sheared pupil, 88 sign convention, 3, 17, 23 Sine condition, 162–164 for the afocal case, 167 in the finite conjugate, 162 with the object at infinity, 163 skew rays, 46–48, 55, 67, 188 Snell's law, 2–4, 39, 45–49,53–54 60, 105, 110, 126 spatial frequency, 67, 86-89, 174, 195, 249 spheres, 8 spherical aberration, 72-75, 85-88, 102-103,112-114, 122-130, 135-137, 142, 148-156, 158, 162, 163, 170-173, 177-178, 182, 184–188, 203, 234–241, 249, 253-259, 262, 265-268, 272, 276 higher-order, 75, 82, 85, 169, 174, 177, 186, 201-202 mirrors, 129 sagittal oblique, 188 surface, 160 tangential oblique, 188 spherochromatism, 100, 177–178, 267 - 268split doublets, 177 splitting lenses, 156 spot diagrams, 69-71, 75-78, 89, 99, 101, 173–175, 177–179, 234–246, 273–275, 278, 281 surface aplanatic, 122–125, 149, 164, 170, 195

aspheric, 57, 120, 128, 157–162, 233-236, 239, 241-242, 253, 256, 258, 259, 261, 262, 263, 264, 266, 268, 269, 271 image, 79, 207 normal to, 2, 60object, 65 paraxial, 70, 86 at reflecting, 61 Petzval, 132–135, 154 reflecting, 3, 18, 130, 254 refracting, 5, 21–23, 48, 57, 102, 162, 255 sagittal, 154 image, 133–134, 184, 188, 207 spherical, 160 tangential, 154 image, 133–134, 188 toroidal, 57

Т

tangential astigmatism, 85, 185 coma, 241 curve, 173, 179, 186-188, 203 foci, 184 focus, 78 image surface, 133-134, 188 flat, 184 plane, 131 section, 67, 75, 77-78, 83, 206 surface, 154 telephoto lens, 158, 193-197, 279-282 telescope objective, 97 thermal imaging lenses, 153 thick lens, 20, 33, 35–36, 92, 155 thin lens, 33, 38, 92, 97, 111, 122, 139–157, 168–169, 177, 183, 189, 193 third-order aberrations, 27, 75–76, 85–89, 102, 106, 112, 159, 170, 175, 202

spherical, 194 astigmatism, 133 distortion, 79, 86 through-focus, 75-78 toroidal surface, 57 transverse aberration, 45–46, 63–68, 77, 83-85, 129-131, 134, 168, 173–175, 180, 185, 188, 190, 193, 197, 203–207 of coma corrector, 260 of field corrector, 258, 261, 267 for a Cassegrain telescope, 258 for a Ritchey-Chrétien telescope, 265 of the achromatized Schmidt camera, 270 of the Canzek Mangin system, 278of the field-flattened Schmidt camera. 271 of the Maksutov-Bouwers Cassegrain system., 273 of the Ritchey-Chrétien telescope, 263 of the Schmidt camera, 269 of the Wiedemann-Mangin mirror system, 275 magnification, 28 trigonometric ray tracing, 47 triplet, 113, 194–195, 199–208

U

undercorrected aberrations, 128 unit magnification, 19–20

V

vignetting, 65, 67, 186, 206–207 V-values of glass, 169

W

wavefront, 5-6, 178, 182

aberration, 45, 55, 63–78, 83–88, 71-77, 101–116, 130–131, 91, 99–100, 146, 159, 165, 262 geometrical, 55 Wiedemann-Mangin, 274–276 Wiedemann-Mangin mirror, 274, 275

Ζ

zero Seidel conditions, 126



Kidger was Michael born in Birmingham, England, on July 6, 1937. He achieved scholarships at the age of 17 to Imperial College, London, where he graduated in 1958, and was awarded an MSc in Applied Optics in 1959. He spent a short time in industry with the optical firm Taylor, Taylor and Hobson of Leicester. In 1963 he joined the optical design team at Imperial College under Professor Charles Wynne. In 1966 he accompanied Wynne to work on bubble chamber optics at the Brookhaven Laboratories. In 1967 Kidger was appointed lecturer in the applied optics section of

Imperial College—a post he held for 20 years. He retained a part-time teaching post at IC and left in 1987 after an association lasting 33 years. His PhD dissertation, "The Application of Electronic Computers to the Design of Optical Systems, Including Aspheric Lenses," was published in 1971. In 1982 he formed the company Kidger Optics Ltd. with his wife Tina. He was a regular participant and exhibitor at SPIE meetings and served on several SPIE committees, including the Scholarship Committee. He gave optical design courses worldwide up until his death. Michael Kidger died in Australia on February 2, 1998, at the age of 60.