

Journal of Biomedical Optics

SPIEDigitalLibrary.org/jbo

Errata: Altered lymphatic function and architecture in salt-induced hypertension assessed by near-infrared fluorescence imaging

Sunkuk Kwon
Germaine D. Agollah
Wenyaw Chan
Eva M. Sevick-Muraca



Errata: Altered lymphatic function and architecture in salt-induced hypertension assessed by near-infrared fluorescence imaging

Sunkuk Kwon,^a Germaine D. Agollah,^{a,b} Wenyaw Chan,^c and Eva M. Sevick-Muraca^a

^aUniversity of Texas Health Science Center, Center for Molecular Imaging, The Brown Foundation Institute of Molecular Medicine, Houston, Texas 77030

^bUniversity of Texas Graduate School of Biomedical Sciences at Houston, University of Texas MD Anderson Cancer Center, Houston, Texas 77030

^cUniversity of Texas Health Science Center at Houston, School of Public Health, Houston, Texas 77030

[DOI: 10.1117/1.JBO.17.8.089802]

This article [*J. Biomed. Opt.* **17**, 080504 (2012)] was originally published online on 16 August 2012 with errors in the caption and callout of Fig. 1. A sentence was removed from the caption, and two references to figure parts were changed in text.

On p. 2, column 2, the first full sentence was changed from:

“In addition, our data showed significantly increased fluorescent intensities in the ILN of HSD-fed mice as compared to baseline, indicating increased lymph flow. Similar to these results in mice, our data also demonstrated dilated lymphatic vessels in the ear of rats fed a HSD [Fig. 1(f)], which were confirmed by immunofluorescence staining of lymphatic vessels [Fig. 1(d) and 1(h)].”

To:

“In addition, our data showed significantly increased fluorescent intensities in the ILN of HSD-fed mice as compared to baseline data in the hindlimb [inset in (a)] and in the ear (d). (c) Significantly increased fluorescent intensities in the ILN of HSD-fed mice were observed as compared to baseline. (f) The apparent average lymphatic vessel diameter associated with HSD was significantly enlarged as compared to baseline. *Inset*, magnified fluorescent images of the dashed rectangles. *Asterisk*, ILN. *Arrow*, ICG injection site. *Scale bar*: 1 mm. Dashed lines in (d) and (e) indicate the boundary of the injection site. * $p < 0.05$ versus baseline.”

Figure 1 and the corrected caption are reprinted here. All versions of the article were corrected on 21 August 2012.




Fig. 1 Dilated lymphatic vessels were observed in the hindlimb (inset in (b)) of HSD-fed mice and in the ear of HSD-fed rats (e) as compared to baseline data in the hindlimb [inset in (a)] and in the ear (d). (c) Significantly increased fluorescent intensities in the ILN of HSD-fed mice were observed as compared to baseline. (f) The apparent average lymphatic vessel diameter associated with HSD was significantly enlarged as compared to baseline. *Inset*, magnified fluorescent images of the dashed rectangles. *Asterisk*, ILN. *Arrow*, ICG injection site. *Scale bar*: 1 mm. Dashed lines in (d) and (e) indicate the boundary of the injection site. * $p < 0.05$ versus baseline.