Errata: Influence of loupes and age on the near visual acuity of practicing dentists

Martina Eichenberger,^a Philippe Perrin,^a Klaus W. Neuhaus,^a Ueli Bringolf,^b and Adrian Lussia^a

^aUniversity of Bern, School of Dental Medicine, Department of Preventive, Restorative and Pediatric Dentistry, CH-3010 Bern, Switzerland

^bNordstrasse 26, CH-8200 Schaffhausen, Switzerland

[DOI: 10.1117/1.3600762]

This article [*J. Biomed. Opt.* **16**, 035003 (2011)] was originally published online on 16 Mar 2011 with an error in Table 4: the columns titled “95% CI” were reversed. The corrected table appears below.

The article was corrected online on 27 May 2011.

Table 4 Visual acuities for the groups with a natural visual acuity ≥ 1.1 and < 1.1 under different magnification conditions: natural visual acuity (NV), single lens loupe (S), Galilean system (G), and Keplerian system (K). Within the columns, significant differences are represented by the different superscript letteres (a: $p < 0.001$, b: $p < 0.05$).

Condition	Visual acuity ≥ 1.1			Visual acuity < 1.1		
	Mean	Median	95% CI	Mean	Median	95% CI
NV ^a	1.33	1.36	(1.28;1.39)	0.97	1.01	(0.94;1.01)
S, 2 $\times$ ^b	1.58	1.31	(1.31;1.56)	1.19	1.31	(1.13;1.31)
G, 2.5 $\times$ ^b	2.66	2.74	(2.37;2.74)	2.13	2.32	(1.99;2.32)
K, 4.3 $\times$ ^b	5.12	4.62	(4.31;5.22)	4.00	4.00	(3.46;4.31)