Large-field-of-view optical elastography using digital image correlation for biological soft tissue investigation (erratum)

Daniel Claus
Marijo Mlikota
Jonathan Geibel
Thomas Reichenbach
Giancarlo Pedrini
Johannes Mischinger
Siegfried Schmauder
Wolfgang Osten

Large-field-of-view optical elastography using digital image correlation for biological soft tissue investigation (erratum)

Daniel Claus,^a Marijo Mlikota,^b Jonathan Geibel,^a Thomas Reichenbach,^a Giancarlo Pedrini,^a Johannes Mischinger,^c Siegfried Schmauder,^b and Wolfgang Osten^a

^aUniversität Stuttgart, Institut für Technische Optik, Stuttgart, Germany

^bUniversität Stuttgart, Institut für Materialprüfung, Werkstoffkunde und Festigkeitslehre, Stuttgart, Germany

^cEberhard Karls Universität Tübingen, Klinik für Urologie, Universitätsklinikum Tübingen, Tübingen, Germany

[DOI: 10.1117/1.JMI.4.2.029801]

This article [*J. Med. Imag.* 4(1), 014505 (2017)] was originally published with the captions for Figs. 8 and 9 transposed. The captions and figures below are correctly matched.

This article was corrected online on 23 May 2017.

Fig. 8 Stress distributions obtained using the hyperelastic Arruda–Boyce model for Abaqus (a) σ_{xx} at the front surface, (b, c, d) 3-D stress distribution along (b) the x -direction σ_{xx} , (c) y -direction σ_{yy} , (d) and z -direction σ_{zz} with indenter positioned on top of inhomogeneity, (e) 3-D stress distribution along z -direction σ_{zz} with indenter positioned in 10 mm distance to inhomogeneity.

Fig. 9 Flow chart and results for obtained displacements fields with and without foreign body, the resulting difference displacement field, the calculate strain field, and corresponding cross-section plot, which compares the difference approach strain field with the conventional results shown in Fig. 7(b).