Journal of Applied Remote Sensing

RemoteSensing.SPIEDigitalLibrary.org

Application of MODIS land surface temperature data: a systematic literature review and analysis

Thanh Noi Phan
Martin Kappas

SPIE.

Thanh Noi Phan, Martin Kappas, "Application of MODIS land surface temperature data: a systematic literature review and analysis," *J. Appl. Remote Sens.* **12**(4), 041501 (2018), doi: 10.1117/1.JRS.12.041501.

Application of MODIS land surface temperature data: a systematic literature review and analysis

Thanh Noi Phan^{a,b,*} and Martin Kappas^a

^aUniversity of Göttingen, Department of Cartography, GIS and Remote Sensing,
Faculty of Geoscience and Geography, Göttingen, Germany

^bVietnam National University of Agriculture, Department of Cartography and Geodesy,
Faculty of Land Management, Hanoi, Vietnam

Abstract. The Moderate Resolution Imaging Spectroradiometer (MODIS) aboard the Terra and Aqua satellites, which provides a very high temporal (four times per day) and spatial (1 km) resolution, has become one of the most important and widely used sensors for a broad range of applications. We analyze 529 articles from 159 journals in the Scopus database from 2009 to 2018 to understand the global and longitudinal trends of MODIS land surface temperature (LST) data applications. The results show that the publications of papers related to MODIS LST data have been steadily rising annually. They spanned 19 subject areas from environmental, agricultural, and biological science to social science and medicine, indicating a wide range of MODIS LST data applications. Among the 159 journals, *Remote Sensing of Environment*, *Remote Sensing*, and the *International Journal of Remote Sensing* published the most articles. The study also showed that urban heat island (UHI), air temperature estimation/mapping (Ta estimation), soil moisture, evapotranspiration estimation, and drought monitoring/estimation were the most popular applications of MODIS LST data. Furthermore, we discuss the strengths, limitations, and future direction of research using MODIS LST data. © 2018 Society of Photo-Optical Instrumentation Engineers (SPIE) [DOI: [10.1117/1.JRS.12.041501](https://doi.org/10.1117/1.JRS.12.041501)]

Keywords: land surface temperature; moderate resolution imaging spectroradiometer land surface temperature products; moderate resolution imaging spectroradiometer land surface temperature applications.

Paper 180278V received Apr. 4, 2018; accepted for publication Sep. 28, 2018; published online Oct. 25, 2018.

1 Introduction

Land surface temperature (LST) is the “skin” temperature of the earth surface, a key parameter for land-surface processing studies at both a regional and global scale.^{1–5} It plays an important role in a wide range of fields, such as environmental monitoring,⁶ hydrology,⁷ urban climate, urban heat island (UHI),^{8–10} and agriculture.^{11–13} Due to rapid changes of LST on spatial and temporal scales, ground-based station observations cannot represent the LST of a region. In the other words, the observations from stations become insufficient for large areas.³ Remotely sensed image data is one of the most suitable sources of LST data at high spatial and/or temporal resolution.^{3,14} However, it is worth noting that, due to the trade-off between spatial and temporal resolution when designing satellite sensors, most of the single satellites cannot provide images with both spatial and temporal high resolution.¹⁵ In general, if a satellite provides high spatial resolution images, then its temporal resolution is medium/low or vice versa.¹⁶ For example, Advanced Very High Resolution Radiometer (AVHRR) and Moderate Resolution Imaging Spectroradiometer (MODIS) can provide daily LST data, but their spatial resolution is about 1000 m; meanwhile, Landsat can provide LST at 100-m resolution, but the temporal resolution is 16 days. Regarding the single satellite sensors, among various remote sensing sensors that provide LST data, the most popular sensors are the MODIS, AVHRR, Landsat (ETM+, OLI), and the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER).

*Address all correspondence to: Thanh Noi Phan, E-mail: thanh-noi.phan@geo.uni-goettingen.de

To produce an overview of LST data applications, we searched {"sensor-name" AND ("land surface temperature" OR "LST")} within titles, abstracts, and keywords in the Scopus database (accessed on July 19, 2018). Articles published in book series were removed from the results so that only articles published in journals and conference proceedings in the English language were chosen. The results show that the number of publications with MODIS was highest when compared with AVHRR, Landsat, and ASTER (Fig. 1). The number of publications relating to MODIS LST has been steadily increasing from 2009 to 2018.

The results in Fig. 1 confirm that MODIS LST data are the most commonly used, which was stated in many previous studies.^{17–19} Similar to MODIS LST data, AVHRR can provide very high temporal resolution (twice daily) and spatial resolution (~1.1 km) data; however, some studies have compared LST from MODIS and AVHRR with *in situ* LST and found that the difference between the *in situ* data and AVHRR was larger than those of MODIS.^{20,21} As shown in Fig. 1, the number of publications using ASTER images has generally remained steady and low. One explanation could be that, although ASTER can provide very high spatial resolution (90 m) and a temporal resolution of 16 days, it is not free of charge and the data are only provided upon request.²² Regarding the Landsat data, similar to MODIS, the number of publications has been on the rise between 2009 and 2018. From 2009 to 2015, the number Landsat publications was much lower than of MODIS; however, over the last 3 years (2016, 2017, and 2018) the number of publications using Landsat data has significantly risen, approaching the number of MODIS data publications (Fig. 1). This rising can be explained that because there has been a significant increase in applications of downscaling techniques, which mainly downscale MODIS LST data to the higher resolution (i.e., Landsat), or use MODIS LST and Landsat LST for downscaling and applied for specific applications.^{23–29}

In this study, MODIS LST data indicate all MODIS LST data available and free for download from NASA Land Data Products and Services (Table 1).

The daily MODIS LST data products, MOD/MYD11A1 at 1 km and MOD/MYD11B1 at 6 km, are retrieved using the generalized split-window³⁰ and day/night algorithms,³¹ respectively. MOD/MYD11C1 data products are derived from the MOD/MYD11B1 product, and stored in a 0.05 deg geographic climate modeling grid (CMG). The 8-day MODIS LST data

Fig. 1 The total number of articles for each related keyword searched in the Scopus database until July 19, 2018.

Table 1 Different types of MODIS LST data used for this review.

Products	Terra	Aqua
Land surface temperature/emissivity daily L3 global 1 km	MOD11A1	MYD11A1
Land surface temperature/emissivity 8-day L3 global 1 km	MOD11A2	MYD11A2
Land surface temperature/emissivity daily L3 global 6 km	MOD11B1	MYD11B1
Land surface temperature/emissivity 8-day L3 global 6 km	MOD11B2	MYD11B2
Land surface temperature/emissivity monthly L3 global 6 km	MOD11B3	MYD11B3
Land surface temperature/emissivity daily global 0.05 deg CMG	MOD11C1	MYD11C1
Land surface temperature/emissivity 8-day global 0.05 deg CMG	MOD11C2	MYD11C2
Land surface temperature/emissivity monthly global 0.05 deg CMG	MOD11C3	MYD11C3
Land surface temperature/emissivity 5-min L2 Swath 1 km	MOD11_L2	MYD11_L2

products, MODIS 11A2, 11B2, and 11C2, are retrieved by averaging from 2 to 8 days of the corresponding daily products. The monthly products, MOD/MYD 11B3, 11C3, are a monthly composite of LST at the same resolutions as 8-day MOD/MYD 11B2, and 11C2 products. MOD/MYD11_L2 products are the LST products at 1-km spatial resolution for a swath, which are produced daily at 5-min increments, using the generalized split-window algorithm.¹⁷

In addition, it is worth mentioning that, together with MOD/MYD11 data products, which were produced over nearly the last two decades, a new product, MOD/MYD21 LST and emissivity product, was planned to be released early in 2017 with MODIS collection 6 (C6) data. However, these products are currently unavailable due to science data quality issues.³² Although MOD/MYD11 use generalized split-window and day/night algorithms, MOD/MYD21 is generated based on the ASTER temperature emissivity separation (TES) algorithm.³³ According to Coll et al.,³⁴ MODIS 21 TES algorithm is highly complementary to the MODIS 11 algorithms as it provides more accurate LST over semiarid and arid areas, but less accurate LST over heavily vegetated areas and water bodies, whereas MODIS 11 has its largest uncertainties in semiarid and arid areas, but optimum performance over heavily vegetated areas and water bodies.

To the best of our knowledge, no previous review article has studied the applications of MODIS LST data in the literature. Therefore, it is practical to review the application of MODIS LST data over the last 10 years. The first objective is to provide a general overview and guidance about the popular applications of MODIS LST for prospective users. We believe this information will support and encourage novice researchers, particularly in developing countries, to study, evaluate, and apply MODIS LST data. The second objective is to analyze the trends and popular topics of MODIS LST related studies and the potential applications of this data.

2 Materials and Methods

2.1 Bibliographic Database

For many years, Web of Science (WoS) from the Thomson Reuters Institute for Scientific Information (ISI) was the only citation and publication database covering all domains of science. However, in 2004, Elsevier Science introduced the Scopus database, which is an effective alternative to WoS according to Vieira and Gomes.³⁵ Studies comparing WoS and Scopus have concluded that, when comparing different subject areas, an advantage in one of these databases is shown in Ref. 36. Studies comparing WoS and Scopus which focused on a specific field of study; however, concluded that it is difficult to determine the better database.^{37,38} In addition, Vieira and Gomes³⁵ stated that Scopus contains a broader range of journals and provides 20% more

coverage than WoS. This finding is consistent with studies by Jacso³⁹ and Falagas et al.,⁴⁰ which confirmed that Scopus covers more journals and analyzes citations faster than WoS. Moreover, Boyle and Sherman⁴¹ recommended that Scopus should be chosen because of its outcome quality, time-saving features, ease of use, and larger amount of search results.

Currently, with almost 23,000 titles and 150,000 books from ~5000 publishers, Scopus is considered one of the largest abstract and citation databases of peer-reviewed research literature.⁴² We used the Scopus database to select publications for this study (accessed on July 12, 2018).

2.2 Methods

We followed four crucial steps for a systematic review outlined by Stewart:⁴³ (1) obtain all publications in the interested topic by using broad search keywords, (2) use strict and clear criteria to limit the universal results to targeted/eligible publications, (3) extract and code essential information from the eligible publications for the statistical outcome measures, and (4) report and discuss the methods, results based on the selected publications (Fig. 2).

As stated in Sec. 2.1, in this study, we used the Scopus database to search for publications. It should be noted that using the same search keywords, there are 1101 publications within the timespan 1999 to 2018; however, most of the studies (991 publications or ~90%) were published in the period from 2009 to 2018. Therefore, we analyzed publications within this timespan.

First, we performed an advanced search in the Scopus database on July 12, 2018 using the keywords {"MODIS" AND ("land surface temperature" OR "LST")} in titles, abstracts, and keywords (including author keywords and index keywords) from 2009 to 2018. This search resulted in 1451 records. Next, we excluded conference papers, book chapters, review articles, letters, notes, and non-English articles, which resulted in 991 peer-reviewed articles (including published articles and articles in press) published in 159 journals. We further filtered this list by removing 136 articles that did not use MODIS LST data. Next, based on the titles, abstracts, and full texts (if necessary), we removed articles that focused more on theories and methods, or only analyzed the spatial and temporal variations of LST. As a result, 529 articles were selected for the final list (Fig. 2).

Fig. 2 The procedure for selecting articles from Scopus database.

3 Results

3.1 General Research Output Based on Selected Publications

In this section, we present an overview of research application trends using MODIS LST data based on the identified 529 articles published in 159 journals from January 2009 to July 2018.

Generally, the number of publications that used MODIS LST data increased from 2009 to 2017. It should be noted that the publications focusing on theories, methodologies, or validation purposes have been removed from this review study. In 2018, by the date of the Scopus search (July 12), there were 55 articles already published (or in press). It is reasonable to predict that this increasing trend will continue through 2018 (Fig. 3).

Regarding the subject classification of the publications, 529 articles were published within 19 subject areas in 159 journals. The top three subject areas were Earth and Planetary Sciences (397 articles—75%), Environmental Science (172 articles—32.5%), and Agricultural and Biological Sciences (164 articles—31%). The total number of articles was not equal to 529 because many belong to multiple subject areas. In addition, the studies of MODIS LST data were published in both general research area journals (e.g., *J-STARS*, *Hydrology*, and *Journal of Earth System Science*) and specific journals (e.g., *Remote Sensing of Environment* and *International Journal of Climatology*), indicating a wide range applications of MODIS LST data.

Table 2 shows the 11 most active journals that have published MODIS LST data-based research. Among the 529 articles, 255 articles (48.2%) were published in these 11 journals, whereas the remaining 274 (51.8%) were published in the other 148 journals. In addition, 65.8% of total citations (TC) were received for these 11 journals, whereas only 34.2% were received for the other 148 journals. The three major publication outlets for MODIS LST data-based research were *Remote Sensing of Environment*, *Remote Sensing*, and *International Journal of Remote Sensing*, with a total number of articles (% articles) of 62 (11.7%), 56 (10.6%), and 30 (5.7%), respectively. Table 2 shows that the journals with the highest TC values were *Remote Sensing of Environment* with 2808 citations (33.5%),

Fig. 3 The total publications (gray—855 articles) and applied for a specific field (orange—529 articles) in Scopus database from 2009 to 2018.

Table 2 The top most active journals publishing research based on MODIS LST data.

Ranking	Journal	Total articles (%)	Total citations (%)	Citations per articles
1	Remote Sensing Of Environment	62 (11.7)	2808 (33.5)	45.3
2	Remote Sensing	56 (10.6)	397 (4.7)	7.1
3	International Journal Of Remote Sensing	30 (5.7)	273 (3.3)	9.1
4	Journal Of Geophysical Research Atmospheres	20 (3.8)	293 (3.5)	14.7
5	ISPRS Journal Of Photogrammetry And Remote Sensing	17 (3.2)	344 (4.1)	20.2
6	International Journal Of Applied Earth Observation And Geoinformation	15 (2.8)	353 (4.2)	23.5
7	IEEE Journal Of Selected Topics In Applied Earth Observations And Remote Sensing (J-STARS)	13 (2.5)	73 (0.9)	5.6
8	Theoretical and Applied Climatology	12 (2.3)	353 (4.2)	29.4
9	Journal of Hydrology	10 (1.9)	215 (2.6)	21.5
9	Agricultural and Forest Meteorology	10 (1.9)	289 (3.4)	28.9
9	Hydrology and Earth System Sciences	10 (1.9)	126 (1.5)	12.6

Table 3 Top 10 most cited articles.

	Articles	Cited by	References
1	Remote sensing of the UHI effect across biomes in the continental USA	344	Imhoff et al. ⁴⁴
2	Evaluation of MODIS land surface temperature data to estimate air temperature in different ecosystems over Africa	199	Vancutsem et al. ⁴⁵
3	Monitoring agricultural drought for arid and humid regions using multisensor remote sensing data	141	Rhee et al. ⁴⁶
4	Downscaling SMOS-derived soil moisture using MODIS visible/infrared data	139	Piles et al. ⁴⁷
5	Estimating air surface temperature in Portugal using MODIS LST data	127	Benali et al. ⁴⁸
6	Estimating volumetric surface moisture content for cropped soils using a soil wetness index based on surface temperature and NDVI	109	Mallick et al. ⁴⁹
7	Exploring indicators for quantifying surface UHIs of European cities with MODIS land surface temperatures	103	Schwarz et al. ⁵⁰
8	Estimation of daily maximum and minimum air temperature using MODIS land surface temperature products	86	Zhu et al. ⁵¹
9	Comparison of multiple models for estimating gross primary production using MODIS and eddy covariance data in Harvard Forest	85	Wu et al. ⁵²
10	Estimation of net radiation from the MODIS data under all sky conditions: Southern Great Plains case study	84	Bisht and Bras ⁵³

Remote Sensing with 397 citations (4.7%), and the *International Journal Of Applied Earth Observation and Geoinformation* with 353 citations (4.2%).

Table 3 shows the 10 most cited articles, which cover five application topics: UHI effect, Ta estimation, drought monitoring, soil moisture, estimating gross primary production, and

net radiation estimation. Among the 10 most cited articles, three articles focus on Ta estimation, two articles focus on UHI effect, and two articles focus on soil moisture applications. Again, it should be noted that the citation count in this review only used the Scopus database (other databases, e.g., Google Scholar, could show a higher number of citations). As shown in Table 3, the most cited article was written by Imhoff et al.⁴⁴ with 344 citations, followed by Vancutsem et al.⁴⁵ with 199 citations, and Rhee et al.⁴⁶ with 141 citations. These articles studied UHI, Ta estimation, and drought monitoring. An interesting finding is that these top three articles were published in the same year (2010), suggesting that these three topic applications could be the most applications of MODIS LST data.

3.2 Main Applications of MODIS LST Data

In order to get an overview of MODIS LST applications, we generated a word cloud using the titles of the 529 selected articles. As shown in Fig. 4, the most commonly used words were urban, heat, estimation, air, soil, island, and moisture. Based on this result and further investigating the titles, abstracts, and full text (if required) of the 529 selected articles, we found that the top five applications were UHI (63 articles—11.9%), air temperature (Ta) estimation (52 articles—9.8%), soil moisture estimation (50 articles—9.5%), evapotranspiration (48 articles—9.1%), and drought (33 articles—6.2%). This is consistent with the hypothesis we have made based on the citations trend shown in Table 3. The remaining (~50%) were applied for various application fields, such as risk mapping (e.g., earthquakes, forest fires, mosquitoes, and floods); agriculture (rice yields, paddy rice mapping, and rice quality); water (surface water temperature, lake temperature, and water pollution); hydrological models; urban expansion; snow melt; snow cover; and many others.^{54–62}

Among these top five applications of MODIS LST, three applications could be considered significantly more popular because more than 50% of publications for these applications were published in the last 3 years. These applications are drought monitoring/assessment (21 articles—63.6%), UHI effect (35 articles—55.6%), and Ta estimation (27 articles—52%). However, in comparison with UHI or Ta estimation studies, drought studies used additional data from MODIS sensors (and other sensors). In other words, MODIS LST is considered a key input for UHI and Ta studies; however, in drought studies, MODIS LST is just one input among many other important inputs (i.e., NDVI from MODIS, precipitation from TRMM).^{46,63–66} Therefore, in the next section, we will discuss the UHI effects and Ta estimation/mapping studies.

Fig. 4 (a) The cloud of the words contained in the titles of 529 selected articles. The size of each word is indicative of its relative frequency of occurrence and (b) the top 25 most frequently used words.

3.3 UHI Using MODIS LST Data

The UHI, a well-known effect of urbanization, is a common phenomenon, where surface temperatures are higher in urban areas than in the surrounding rural areas.⁶⁷ This effect is considered one of the most significant impacts of human activity on the Earth's surface climate.^{68,69} Although UHI are localized hotspots in the landscape, they have a profound impact on the lives of the urban residents, who currently account for more than half of the total population; urban population is expected to rise about 66% by 2050.^{70,71} In 2014, the number of people living in urban areas in developed regions such as Northern America, Latin America and the Caribbean, and Europe was estimated to be ~82%, 80%, and 73%, respectively. Meanwhile, the urban population of developing regions was around 48% and 40% for Asia and Africa, respectively. The UN⁷⁰ further predicted that the fastest urbanization will take place in Asia and Africa, with urban population predictions of about 64% and 54% by 2050, respectively.⁷⁰ This indicates that in these two regions, the urban landscape will change significantly, and therefore resources should be focused there to implement well-planned and sustainable development.

Traditionally, UHI studies were based on the data retrieved from ground weather stations.^{72,73} Some limitations of UHI studies using weather station data have been reported in the literature because: (1) the distribution of weather stations, particularly in developing countries or remote regions, is usually sparse; therefore, it cannot represent the UHI for a whole city,⁷⁴ (2) most weather stations are located in urban areas, so rural station data for UHI analysis is lacking,⁷⁵ and (3) quality data from stations are usually not available to all users.⁷³ In contrast, remote sensing satellites provide easy to access LST data with global coverage at no cost and can therefore be considered the most suitable way to investigate the surface UHI (SUHI).

Rao⁷⁶ first reported the perspective of using remote sensing techniques for SUHI studies. In the subsequent years, many scholars stated that, in UHI studies, satellite temperature measurement produced better results than interpolating weather station observations.^{77,78} Since the 1990s, the AVHRR sensor has been widely used for LST analysis^{79,80} and the SUHI effect at the regional scale.^{81,82} Later, ASTER, which provides both LST daytime and nighttime data at high spatial resolution (90 m), has been applied for SUHI studies in many cities.^{83–85} Deilami et al.⁸⁶ produced a systematic review of spatiotemporal factors, data, methods, and mitigation measures of the UHI effect, investigating all English peer-reviewed articles published from January 1965 to July 2017. The results showed that the most widely used LST data for Ta were Landsat TM (54.6%), followed by Landsat ETM (34.6%) and MODIS (28.0%). The other satellite images used for UHI studies were Landsat 8,^{87–90} SPOT,⁹¹ and QuickBird.⁹² However, as noticed by the USGS in August 2013, surface water temperatures derived from Landsat 8 TIRS data may be inaccurate, potentially impacting the results of research.⁹³ Obviously, this calibration problem has restricted the application of Landsat 8 TIRS for UHI studies.⁹⁴

Although Landsat is considered one of the most popular satellite data sources for UHI studies, according to Deilami et al.,⁸⁶ there are two main shortcomings of Landsat data: (1) the low temporal resolution (16 days) is not suitable for diurnal or weekly UHI monitoring or assessment and (2) due to the small size of Landsat images, multiple images must be mosaiced when applied to regional and national areas. According to Irons et al.⁹⁵ and Wulder et al.,⁹⁶ the mosaic processing takes a significant amount of time. Currently, it is worth noting that the Landsat surface reflectance level-2 products can be retrieved from the NASA website at no cost, but the preprocessing time can be large when requesting multiple images for a large area.

To overcome the limitations of Landsat satellite data over the last decade, MODIS data have been widely applied to SUHI studies, particularly to those focusing on temporal variation, i.e., diurnal, seasonal, annual, decadal, and coarser scales.^{50,97–99} In addition, it is worth noting that when investigating the SUHI using remote sensing LST data, not only the surface characteristics should be taken into consideration but also the climatic and meteorological conditions.^{100,101} The potential applications and the rising trend of using MODIS data have been reported by Zhang et al.,¹⁰² who conducted a bibliometric analysis of global remote sensing research from 2010 until 2015. To trace the temporal trend of the main applications using remote sensing techniques, they investigated the top keyword from each year (from 2010 to 2015), and the results showed that MODIS emerged as the core theme. The “MODIS” keyword ranked in the top six keywords (among “model,” “classification,” “algorithm,” “SAR,” and “remote sensing”).

As mentioned in Sec. 3.2, one of the most popular applications of MODIS LST data is for UHI/SUHI studies, 12.3% (65 articles). It is worth noting that in UHI/SUHI studies, LST data from Landsat have been widely applied.⁸⁶ However, in recent years, the UHI studies are transitioning from single date to time series (multiple day), from local and regional to global. In addition, the variation of UHI/SUHI due to the changing seasons, daytime and nighttime, has been receiving the attention from scholars. Therefore, remote sensing image data that provide high temporal resolution (i.e., daily, hourly) and coverage of large areas (i.e., city) is becoming indispensable. MODIS satellite data, which provide four LST datasets each day at 1-km spatial resolution with global coverage, are among the most suitable sources for UHI studies at the regional and global scales. Another advantage of MODIS satellite data is that it provides readily available products for UHI applications (e.g., LST and land cover data), which significantly reduces the processing time. Among the 65 articles using MODIS LST data for SUHI studies, 13 articles were focused on analyzing diurnal SUHI, 19 articles analyzed the effects of season on SUHI intensity, and 37 articles investigated the effects of daytime and nighttime on SUHI intensity. Among these 37 articles, two studies (Haashemi et al.⁸ and Bahi et al.¹⁰³) used nighttime LST data of MODIS and daytime LST from Landsat images. It should be noted that among the 65 articles, the numbers of studies that were conducted in China, India, and the United States were 21, 9, and 7, respectively. The remaining articles were mostly implemented in EU cities. However, based on the prediction of UN,⁷⁰ urbanization will take place in developing countries regions, Africa, and Asia; this suggests that MODIS LST will be one of the most valuable data sources for the hot topic, UHI studies, in developing countries in the near future.

Another important point is that there are four MODIS LST data images at four times each day. Between the two satellites (Terra and Aqua) collecting these data, the overpass time of the Aqua satellite (1:30 a.m. and 1:30 p.m.) is closer to the occurrence of the highest and lowest temperatures in the diurnal cycle than the Terra satellite (10:30 a.m. and 10:30 p.m.). However, most studies choose to utilize either Terra data^{9,103–107} or Aqua data^{44,73,108–110} without explanation. According to Clinton and Gong,¹¹¹ the daytime SUHI monitored by Aqua LST (1:30 p.m.) was higher than that of Terra (10:30 a.m.). Therefore, a topic, for example, the effects of different time observations on UHI, should be taken into consideration for future studies.

3.4 Air Surface Temperature Estimation Using MODIS LST Data

Air surface temperature (T_a), which is usually measured at weather stations 2 m above the ground, is a key parameter for a wide range of applications.^{48,112–114} Although it provides accurate T_a , it has limitations in spatial coverage. Traditionally, to obtain T_a for a large area, interpolation methods were used,^{115,116} however, the accuracy of interpolation methods strongly depends on the distribution of weather stations.^{45,117} Therefore, it is difficult to apply this method for regions with sparse weather stations, such as in developing countries and remote areas.

According to Li et al.²⁴ and Chen et al.,¹¹⁸ remotely sensed LST data are considered one of the most suitable ways to retrieve T_a data at local, regional, or global scales. Among the remote sensing satellites, the Advanced Very High Resolution Radiometer (AVHRR), Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER), and Landsat can provide LST data, though LST data from MODIS are the most widely used.¹⁹ To confirm this, we used the same search terms for ASTER, AVHRR, and Landsat (which resulted in 52 publication using MODIS LST). The result showed that from 2009 to 2018 there were only four records total; three for Landsat and one for ASTER.^{119–122} This result indicates that MODIS LST is the most popular and widely used source for T_a estimation studies. Among these four articles, Zhu et al.¹²² used both Landsat 8 and MODIS LST for T_a estimation, showing that T_a could be estimated at the 30-m (using Landsat 8) and 1000-m scales (using MODIS) with a root mean square error (RMSE) of 1.7°C and 2.6°C, respectively. Among the 52 T_a estimation publications collected for this study, the number of studies that applied MODIS LST for daily, weekly, and monthly T_a estimation were 45, 2, and 7, respectively. The total number is not equal to 52 because two studies were applied for both daily and weekly T_a ,^{48,123} and one study was applied to estimate both daily and monthly T_a .¹²⁴ In addition, the difference between daily, weekly, and monthly articles also suggests that the high temporal resolution of MODIS LST data (four times daily) is one of the largest advantages of MODIS data and is, therefore, widely applied for T_a estimation.

Three common method types were used: statistical methods (78.8%—41 articles), machine learning methods (13.5%—7 articles), and temperature vegetation index methods (7.7%—4 articles). It is worth noting that in most of the articles using statistical or machine learning methods, MODIS LST data were combined with auxiliary variables. The most widely used auxiliary variables were elevation, NDVI, Julian day, latitude, longitude, and day length.^{123,125–127} Although statistical methods were the most widely used, by 2013 the researchers started focusing on machine learning approaches.^{127–133} Some studies compared the performance of advanced statistical methods and machine learning methods but with inconclusive results. Xu et al.¹²⁹ compared two methods, linear regression, and random forest (RF) for Ta estimation in British Columbia, Canada, concluding that the RF model achieved better accuracy [mean absolute error (MAE) = 2.02°C, $R^2 = 0.74$, compared with MAE = 2.41°C, $R^2 = 0.64$]. Meyer et al.¹³⁰ compared a simple linear regression model with three machine learning algorithms [RF, generalized boosted regression models (GBM), and Cubist (CB)] for Ta estimation in Antarctica. Their results showed that machine learning algorithms only slightly outperformed the simple linear estimation model. Among the three machine learning methods, GBM showed the best results for Ta estimation. Zhang et al.¹³¹ compared six statistical models (i.e., multiple linear regression, the partial least squares regression, back propagation neural network, support vector regression, RF, and CB) with two types of MODIS LST data qualities (all clear sky data and only good data). They concluded that the performance of each algorithm depended on the different combinations of MODIS LST and the quality of data. Overall, however, RF and CB showed the best results of Ta estimation with both conditions of data quality. Noi et al.¹³² compared the performance of a linear regression model (LM) with RF and CB algorithms, for two types of data input, MODIS LST data solely and MODIS LST with auxiliary variables (elevation and Julian day). Their results showed that both CB and LM performed well with MODIS LST solely, but when MODIS LST is used with auxiliary variables, CB and RF are recommended. Consistent with Zhang et al.¹³¹ and Noi et al.,¹³² Xu et al.¹³³ compared 10 machine learning algorithms for Ta monthly mapping in the Tibetan Plateau and reported that CB outperformed the other methods.

Regarding the MODIS LST data for Ta estimation, four MODIS LST data are available each day (Terra day, Terra night, Aqua day, and Aqua night), with overpass at local times 10:30 a.m., 10:30 p.m., 1:30 p.m., and 1:30 a.m., respectively. However, most of the studies have chosen only one or two MODIS LST data (among the four available) for Ta estimation. Generally, the selection was based on the overpass time of MODIS LST data. For example, Xu et al.¹²⁹ chose LST of Aqua daytime for Ta-max estimation in British Columbia, Canada; Zhang et al.¹³¹ chose LST of Aqua daytime and Terra nighttime, for Ta-max and Ta-min estimation, respectively; and Yang et al.¹³⁴ chose LST of Aqua daytime and nighttime, for Ta-max and Ta-min estimation, respectively. Recently, some studies have chosen all four MODIS LST data for Ta estimation.^{131,132,135,136}

Most recently, Yoo et al.¹²⁷ proposed a new method for Ta estimation by considering the lag time of heat transport from ground level to 2 m. Therefore, they also used the four LST data taken from the day before, so that they used eight MODIS LST datasets total for daily Ta estimation. They implemented this new method in two megacities, Los Angeles, USA, and Seoul, South Korea, and concluded that LST of the day before were crucial for estimating daily Ta in urban climates.

4 Discussions

It is known that remote sensing is the most suitable data for Earth surface-related studies due to its usefulness for monitoring the Earth surface, analyzing trends and patterns, and forecasting. Among various applications of remotely sensed data, land surface temperature (e.g., MODIS LST) is one of the most important variables for climate research and global change.²⁴ Zhang et al.¹⁰² conducted a bibliometric analysis of global remote sensing research during the period 2010 through 2015, showing that MODIS is the most prevalent data source among all the other remotely sensed data (e.g., Landsat, AVHRR, ASTER, SPOT, and IKONOS).

As mentioned in the methods section (Sec. 2.2), although MODIS LST data have been available since 2000, most studies (~90%) using MODIS LST were implemented in the last 10 years

(2009 to 2018). This indicates that MODIS LST data are receiving a large amount of attention from scholars. In addition, in 2017, all MODIS LST data have been reprocessed by NASA and made available for download at no cost, in a collection called MODIS LST Collection 6 (C6).¹⁹ Fixes and improvements have been made in C6, which were reported in detail in Wan.¹⁷ Of these, removing the cloud-contaminated LST pixels is one of the best advantages of C6 compared with the previous collections (C4.1 and C5). Again, the availability of C6 cannot be understated, as the number of publications using MODIS LST data (all publications, including theories and methods publications) in 2017 was higher than in any previous year (Fig. 3) perhaps as a result. However, in addition to the advantages, MODIS LST data have some shortcomings that should be noted. First, similar to other optical (thermal) remotely sensed images, the most concerning problem of MODIS LST data is the lack of cloud-free images. If clouds are present, MODIS LST products are not available.¹³⁷ In addition, pixels that are partially covered or contaminated by clouds, which were not removed from the MODIS LST data may be present, because they could not be detected by the cloud removing mask algorithms.^{138,139} Obviously, in practice, these pixels may affect study results. Second, the spatial resolution of MODIS LST data is too coarse for some applications, such as agriculture, drought, water management, vegetation phenology,^{140,141} or detailed UHI.⁸⁶ Therefore, developing techniques to retrieve high spatial and high temporal resolution from currently available remotely sensed data is a current area of interest.^{16,142} Third, most of the studies in the literature are using MODIS LST without concern for the quality of MODIS LST. Researchers should play more attention to this issue.

4.1 Reconstruction of MODIS LST Data

Regarding the reconstruction techniques, among the 855 articles (included articles focused on theories or methods developments/assessments), there were only 15 articles that considered missing MODIS LST data. Meanwhile, according to Zeng et al.,¹⁴³ completed cloud free MODIS LST data are rare, particularly during rainy seasons and in humid regions. Furthermore, according to Jin and Dickinson,¹⁴⁴ each day, the cloudy-sky condition is present over more than half the globe. This means that MODIS LST data, which have excluded all the cloudy-sky pixels from $M * D11$ products, lose more than half of LST data time series.¹⁴⁵ In addition, Østby et al.¹⁴⁶ compared 8 years of MODIS LST data with *in situ* observations in Austfonna (an Arctic ice cap located on Svalbard), showing that only 26% of the MODIS LST data are retrieved under actual clear sky conditions, of which the percentages of MODIS LST data under clear sky conditions for winter and summer are about 40% and 20%, respectively. As a result, the subsequent applications that use MODIS LST data will be greatly affected.¹²⁷

Furthermore, many previous studies applied MODIS LST data without any consideration for cloud cover, just simply selecting the measurements under clear sky conditions (using the QC data file) for analysis.^{48,125,131,147}

Among the 15 studies that implemented the reconstruction of MODIS LST data, most only focused on the method development and assessment.^{125,148,149} Only a few studies used this technique for specific applications, e.g., Neteler¹⁵⁰ reconstructed MODIS LST for Ta estimation in a mountainous area. In future research, more studies should investigate the effects, and solutions to solve this problem.

4.2 Downscaling Using MODIS LST Data

Among the 855 articles (including articles focusing on theories and methods), there were 60 articles that applied downscaled methods. Among these, there were three studies on UHI effect,^{151–153} 11 articles for monitoring/estimation evapotranspiration, and 12 articles for soil moisture applications, whereas the remaining articles mainly focused on theories or method developments/assessments.

Three types of downscaling studies used MODIS LST data: (1) downscaling MODIS LST data to higher spatial resolution (e.g., 250 and 30 m), (2) downscaling MODIS LST data to higher temporal resolution data (e.g., using hourly GOES—LST data), and (3) downscaling other lower spatial resolution products (e.g., SMOS: Soil Moisture and Ocean Salinity, AMSR-E: Advanced Microwave Scanning Radiometer—Earth Observing System).

As LST is considered one of the most crucial parameters for a wide range of applications, at various scales, high spatiotemporal resolution LST data are required. Currently, a number of sensors can provide high spatial resolution (i.e., Landsat, ASTER) or high temporal resolution (e.g., MODIS, GEOS), yet no sensor can provide LST data at both high spatial and temporal resolution. Therefore, downscaling (or fusion) techniques play a crucial role in retrieving high spatial and temporal resolution for many applications. Although Landsat images provide very high spatial resolutions (up to 60 m), the temporal resolution is too coarse (16 days); therefore, MODIS LST are normally downscaled to Landsat resolution.^{16,24–26,154}

Another important point is that in developing countries, the agricultural field size and other land uses are small, and indeed much smaller than the spatial resolution of MODIS LST data. Therefore, one pixel contains many land cover types, receiving a mixed signal. To reduce the effect of this issue, downscaling to finer resolutions is necessary.

4.3 Quality of MODIS LST Data

According to Colditz et al.,¹⁵⁵ for a long-term time series, data quality is considered a critical issue. MODIS LST data are not available for locations (pixels) that are covered by cloud.¹³⁷ In addition, Wan¹³⁷ also noted that in MODIS LST products, pixels are still present which are covered by thin clouds that cloud-detecting algorithms cannot identify and remove. Therefore, the results of all applications using MODIS LST data are affected. Ackerman et al.¹³⁸ reported that there are ~15% of contamination pixels existent in MODIS LST products. In addition, Williamson et al.¹³⁹ used ground-based meteorological station observations to evaluate the cloud contamination in clear sky MODIS Terra daytime LST in the southwest Yukon area, showing that 13% to 17% of MODIS LST data were unidentified clouds. This will obviously affect the results of subsequent MODIS LST data applications. However, to date, most studies have not considered this problem. Therefore, further studies should investigate the effect on MODIS LST quality and which cloud detection methods are needed.

5 Conclusions

This review gives an overview as well as some specific applications of MODIS LST data from Terra and Aqua satellites over the last 10 years. In general, the number of publications in peer-reviewed journals using MODIS LST data increases annually, indicating that this source of data is becoming more popular and widely used. Among 19 subject areas with several topics of application, the top five applications over the last 10 years were: UHI, air temperature estimation, soil moisture estimation, evapotranspiration estimation, and drought studies. In addition, in the last 3 years, a high number of publications were focused on three topics: drought monitoring/assessment, UHI, and Ta estimation. This reveals the wide range of applications of MODIS LST data. In spite of the advantages of MODIS LST data, some limitations remain, such as the cloud cover effect (included missing pixels, as well as the pixels covered by thin cloud) and the use of the MODIS data quality file (QC file). Some techniques like downscaling (fusion) and reconstruction will bring opportunities for recent and future applications. This study not only clarifies the global overview of MODIS LST data applications to novice researchers but also provides current research hotspots and future potential applications related to MODIS LST data.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the authors.

Acknowledgments

We would like to acknowledge the Vietnamese government for its financial support. We would like to thank Michael Schlund (Institute of Geography, University of Göttingen) for constructive criticism of the initial version of this paper. We also thank the anonymous reviewers for their valuable comments, which improved our paper.

References

1. J. M. Anderson et al., "A thermal-based remote sensing technique for routine mapping of land-surface carbon, water and energy fluxes from field to regional scales," *Remote Sens. Environ.* **112**, 4227–4241 (2008).
2. A. Karnieli et al., "Use of NDVI and land surface temperature for drought assessment: merits and limitations," *J. Clim.* **23**, 618–633 (2010).
3. H. Shwetha and D. N. Kumar, "Prediction of high spatio-temporal resolution land surface temperature under cloudy conditions using microwave vegetation index and ANN," *ISPRS J. Photogramm. Remote Sens.* **117**, 40–55 (2016).
4. S. Khandelwal et al., "Assessment of land surface temperature variation due to change in elevation of area surrounding Jaipur, India," *Egypt. J. Remote Sens. Space Sci.* **21**, 87–94 (2017).
5. M. Norman and F. Becker, "Terminology in thermal infrared remote sensing of natural surfaces," *Agric. For. Meteorol.* **77**, 153–166 (1995).
6. M. Q. Wu et al., "Generating daily high spatial land surface temperatures by combining ASTER and MODIS land surface temperature products for environmental process monitoring," *Environ. Sci. Process. Impacts* **17**, 1396–1404 (2015).
7. A. N. French and A. Inamdar, "Land cover characterization for hydrological modelling using thermal infrared emissivities," *Int. J. Remote Sens.* **31**, 3867–3883 (2010).
8. S. Haashemi et al., "Seasonal variations of the surface urban heat island in a semi-arid city," *Remote Sens.* **8**, 352 (2016).
9. R. Yao et al., "Temporal trends of surface urban heat islands and associated determinants in major Chinese cities," *Sci. Total Environ.* **609**, 742–754 (2017).
10. H. Li et al., "A new method to quantify surface urban heat island intensity," *Sci. Total Environ.* **624**, 262–272 (2018).
11. N. Patel et al., "Analysis of agricultural drought using vegetation temperature condition index (VTCI) from TERRA/MODIS satellite data," *Environ. Monit. Assess.* **184**(12), 7153–7163 (2012).
12. J. Lessel, A. Sweeney, and P. Ceccato, "An agricultural drought severity index using quasi-climatological anomalies of remotely sensed data," *Int. J. Remote Sens.* **37**, 913–925 (2016).
13. K. Hazaymeh and Q. K. Hassan, "A remote sensing-based agricultural drought indicator and its implementation over a semi-arid region, Jordan," *J. Arid Land.* **9**(3), 319–330 (2017).
14. Z. Li et al., "Satellite-derived land surface temperature: current status and perspectives," *Remote Sens. Environ.* **131**, 14–37 (2013).
15. X. Zhu et al., "Spatiotemporal fusion of multisource remote sensing data: literature survey, taxonomy, principles, applications, and future directions," *Remote Sens.* **10**, 527 (2018).
16. Q. Weng, P. Fu, and F. Gao, "Generating daily land surface temperature at Landsat resolution by fusing Landsat and MODIS data," *Remote Sens. Environ.* **145**, 55–67 (2014).
17. Z. Wan, "New refinements and validation of the collection-6 MODIS land-surface temperature/emissivity product," *Remote Sens. Environ.* **140**, 36–45 (2014).
18. J. Kang et al., "Reconstruction of MODIS land surface temperature products based on multi-temporal information," *Remote Sens.* **10**, 1112 (2018).
19. M. Metz, V. Andreo, and M. Neteler, "A new fully gap-free time series of land surface temperature from MODIS LST data," *Remote Sens.* **9**, 1333 (2017).
20. H. Yang and Z. Yang, "A modified land surface temperature split window retrieval algorithm and its applications over China," *Global Planet. Change* **52** (1), 207–215 (2006).
21. L. Zhong et al., "Estimation of land surface temperature over the Tibetan Plateau using AVHRR and MODIS data," *Adv. Atmos. Sci.* **27**, 1110–1118 (2010).
22. C. J. Tomlinson et al., "Remote sensing land surface temperature for meteorology and climatology: a review," *Meteorol. Appl.* **18**, 296–306 (2011).
23. R. Eswar, M. Sekhar, and B. K. Bhattacharya, "Disaggregation of LST over India: comparative analysis of different vegetation indices," *Int. J. Remote Sens.* **37**(5), 1035–1054 (2016).

24. M. Bisquert, J. M. Sánchez, and V. Caselles, "Evaluation of disaggregation methods for downscaling MODIS land surface temperature to Landsat spatial resolution in Barrax test site," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **9**(4), 1430–1438 (2016).
25. S. Bonafoni, "Downscaling of Landsat and MODIS land surface temperature over the heterogeneous urban area of Milan," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **9**(5), 2019–2027 (2016).
26. Y. J. Choe and J. H. Yom, "Downscaling of MODIS land surface temperature to LANDSAT scale using multi-layer perceptron," *J. Korean Soc. Surv. Geod. Photogramm. Cartogr.* **35**(4), 313–318 (2017).
27. W. Essa et al., "Improved DisTrad for downscaling thermal MODIS imagery over urban areas," *Remote Sens.* **9**(12), 1243 (2017).
28. C. Xu et al., "Downscaling of surface soil moisture retrieval by combining MODIS/Landsat and in situ measurements," *Remote Sens.* **10**(2), 210 (2018).
29. X. Pan et al., "Applicability of downscaling land surface temperature by using normalized difference sand index," *Sci. Rep.* **8**(1), 9530 (2018).
30. Z. Wan and J. Dozier, "A generalized split-window algorithm for retrieving land-surface temperature from space," *IEEE Trans. Geosci. Remote Sens.* **34**(4), 892–905 (1996).
31. Z. Wan and Z.-L. Li, "A physics-based algorithm for retrieving land-surface emissivity and temperature from EOS/MODIS data," *IEEE Trans. Geosci. Remote Sens.* **35**(4), 980–996 (1997).
32. G. Hulley and S. Hook, "MOD21 MODIS/Terra Land Surface Temperature/3-Band Emissivity 5-Min L2 1km V006 [Data set]," NASA EOSDIS Land Processes DAAC, https://lpdaac.usgs.gov/dataset_discovery/modis/modis_products_table/mod21_v006 (July 2018).
33. A. R. Gillespie et al., "A temperature and emissivity separation algorithm for advanced spaceborne thermal emission and reflection radiometer ASTER images," *IEEE Trans. Geosci. Remote Sens.* **36**, 1113–1126 (1998).
34. C. Coll et al., "Test of the MODIS land surface temperature and emissivity separation algorithm with ground measurements over a rice paddy," *IEEE Trans. Geosci. Remote Sens.* **54**, 3061–3069 (2016).
35. E. S. Vieira and J. A. N. F. Gomes, "A comparison of Scopus and Web of Science for a typical university," *Scientometrics* **81**, 587–600 (2009).
36. J. Li et al., "Citation analysis: comparison of Web of Science, Scopus, Scifinder, and Google Scholar," *J. Electron. Resour. Med. Libr.* **7**(3), 196–217 (2010).
37. C. López-Illescas, F. Moya-Anegón, and H. F. Moed, "Coverage and citation impact of oncological journals in the Web of Science and Scopus," *J. Informetrics* **2**(4), 304–316 (2008).
38. A. A. Chadegani et al., "A comparison between two main academic literature collections: Web of Science and Scopus databases," *Asian Soc. Sci.* **9**(5), 18–26 (2013).
39. P. Jacso, "As we may search—comparison of major features of the Web of Science, Scopus, and Google Scholar citation-based and citation-enhanced databases," *Curr. Sci.* **89**(9), 1537–1547 (2005).
40. M. E. Falagas et al., "Comparison of SCImago journal rank indicator with journal impact factor," *FASEB J.* **22**(8), 2623–2628 (2008).
41. F. Boyle and D. Sherman, "Scopus: the product and its development," *Ser. Libr.* **49**(3), 147–153 (2006).
42. Scopus database, "Scopus content at a glance," <https://www.elsevier.com/solutions/scopus/how-scopus-works/content> (July 2018).
43. I. D. Stewart, "A systematic review and scientific critique of methodology in modern urban heat island literature," *Int. J. Climatol.* **31**, 200–217 (2011).
44. M. L. Imhoff et al., "Remote sensing of the urban heat island effect across biomes in the continental USA," *Remote Sens. Environ.* **114**(3), 504–513 (2010).
45. C. Vancutsem et al., "Evaluation of MODIS land surface temperature data to estimate air temperature in different ecosystems over Africa," *Remote Sens. Environ.* **114**(2), 449–465 (2010).

46. J. Rhee, J. Im, and G. J. Carbone, "Monitoring agricultural drought for arid and humid regions using multi-sensor remote sensing data," *Remote Sens. Environ.* **114**(12), 2875–2887 (2010).
47. M. Piles et al., "Downscaling SMOS-derived soil moisture using MODIS visible/infrared data," *IEEE Trans. Geosci. Remote Sens.* **49**(9), 3156–3166 (2011).
48. A. Benali et al., "Estimating air surface temperature in Portugal using MODIS LST data," *Remote Sens. Environ.* **124**, 108–121 (2012).
49. K. Mallick, B. K. Bhattacharya, and N. K. Patel, "Estimating volumetric surface moisture content for cropped soils using a soil wetness index based on surface temperature and NDVI," *Agric. For. Meteorol.* **149**(8), 1327–1342 (2009).
50. N. Schwarz, S. Lautenbach, and R. Seppelt, "Exploring indicators for quantifying surface urban heat islands of European cities with MODIS land surface temperatures," *Remote Sens. Environ.* **115**(12), 3175–3186 (2011).
51. W. Zhu, A. Lu, and S. Jia, "Estimation of daily maximum and minimum air temperature using MODIS land surface temperature products," *Remote Sens. Environ.* **130**, 62–73 (2013).
52. C. Wu et al., "Comparison of multiple models for estimating gross primary production using MODIS and eddy covariance data in Harvard Forest," *Remote Sens. Environ.* **114**(12), 2925–2939 (2010).
53. G. Bisht and R. L. Bras, "Estimation of net radiation from the MODIS data under all sky conditions: Southern Great Plains case study," *Remote Sens. Environ.* **114**(7), 1522–1534 (2010).
54. M. Zoran, "MODIS and NOAA-AVHRR L and surface temperature data detect a thermal anomaly preceding the 11 March 2011 Tohoku earthquake," *Int. J. Remote Sens.* **33**(21), 6805–6817 (2012).
55. M. Bisquert et al., "Application of artificial neural networks and logistic regression to the prediction of forest fire danger in Galicia using MODIS data," *Int. J. Wildland Fire* **21**(8), 1025–1029 (2012).
56. M. J. Dantur Juri et al., "Satellite-derived NDVI, LST, and climatic factors driving the distribution and abundance of Anopheles mosquitoes in a former malarious area in northwest Argentina," *J. Vector Ecol.* **40**(1), 36–45 (2015).
57. D. M. Johnson, "An assessment of pre- and within-season remotely sensed variables for forecasting corn and soybean yields in the United States," *Remote Sens. Environ.* **141**, 116–128 (2014).
58. R. Kimura, M. Moriyama, and S. Bandou, "The relationship between land-surface temperature and rice quality in Tottori prefecture, Japan," *Int. J. Remote Sens.* **36**(22), 5690–5706 (2015).
59. E. T. Crosman and J. D. Horel, "MODIS-derived surface temperature of the Great Salt Lake," *Remote Sens. Environ.* **113**(1), 73–81 (2009).
60. K. Song et al., "Spatiotemporal variations of lake surface temperature across the Tibetan Plateau using MODIS LST product," *Remote Sens.* **8**(10), 854 (2016).
61. L. S. Kuchment et al., "Use of satellite-derived data for characterization of snow cover and simulation of snowmelt runoff through a distributed physically based model of runoff generation," *Hydrol. Earth Syst. Sci.* **14**(2), 339–350 (2010).
62. S. D. Chakraborty, Y. Kant, and D. Mitra, "Assessment of land surface temperature and heat fluxes over Delhi using remote sensing data," *J. Environ. Manage.* **148**, 143–152 (2015).
63. N. T. Son et al., "Monitoring agricultural drought in the lower mekong basin using MODIS NDVI and land surface temperature data," *Int. J. Appl. Earth Obs. Geoinf.* **18**(1), 417–427 (2012).
64. C. Hao, J. Zhang, and F. Yao, "Combination of multi-sensor remote sensing data for drought monitoring over Southwest China," *Int. J. Appl. Earth Obs. Geoinf.* **35**, 270–283 (2015).
65. S. Park et al., "Drought assessment and monitoring through blending of multi-sensor indices using machine learning approaches for different climate regions," *Agric. For. Meteorol.* **216**, 157–169 (2016).

66. L. Zhang et al., "Studying drought phenomena in the Continental United States in 2011 and 2012 using various drought indices," *Remote Sens. Environ.* **190**, 96–106 (2017).
67. L. Rose and M. D. Devadas, "Analysis of land surface temperature and land use/land cover types using remote sensing imagery—a case in Chennai city, India," in *Proc. of the Seventh Int. Conf. on Urban Climate*, Yokohama, Japan, Vol. **29** (2009).
68. E. Kalnay and M. Cai, "Impact of urbanization and land-use change on climate," *Nature* **423**, 528–531 (2003).
69. L. M. Zhou et al., "Evidence for a significant urbanization effect on climate in China," *Proc. Natl Acad. Sci. U.S.A.* **101**, 9540–9544 (2004).
70. United Nations, *World Urbanization Prospects: The 2014 Revision: Highlights*, United Nations, Department of Economic and Social Affairs, Population Division, New York (2015).
71. M. Ranagalage, R. C. Estoque, and Y. Murayama, "An urban heat island study of the Colombo Metropolitan Area, Sri Lanka, based on Landsat Data (1997-2017)," *ISPRS Int. J. Geo-Inf.* **6**, 189 (2017).
72. S. R. Gaffin et al., "Variations in New York city's urban heat island strength over time and space," *Theor. Appl. Climatol.* **94**(1–2), 1–11 (2008).
73. J. Wang et al., "Spatiotemporal variation in surface urban heat island intensity and associated determinants across major Chinese cities," *Remote Sens.* **7**(4), 3670–3689 (2015).
74. D. Zhou et al., "Surface urban heat island in China's 32 major cities: spatial patterns and drivers," *Remote Sens. Environ.* **152**, 51–61 (2014).
75. F. Wang et al., "A new estimation of urbanization's contribution to the warming trend in China," *J. Clim.* **28**(22), 8923–8938 (2015).
76. P. Rao, "Remote sensing of urban heat islands from an environmental satellite," *Bull. Am. Meteorol. Soc.* **53**, 647–648 (1972).
77. P. Dash et al., "Land surface temperature and emissivity estimation from passive sensor data: theory and practice-current trends," *Int. J. Remote Sens.* **23**(13), 2563–2594 (2002).
78. M. Jin, J. M. Shepherd, and C. Peters-Lidard, "Development of a parameterization for simulating the urban temperature hazard using satellite observations in climate model," *Nat. Hazards* **43**(2), 257–271 (2007).
79. C. Ottlé and D. Vidal-Madjar, "Estimation of land surface temperature with NOAA9 data," *Remote Sens. Environ.* **40**, 27–41 (1992).
80. G. G. Gutman, "Multi-annual time series of AVHRR-derived land surface temperature," *Adv. Space Res.* **14**, 27–30 (1994).
81. M. J. L. Garcia et al., "NOAA-AVHRR contribution to the analysis of urban heat islands," in *Proc. 5th Int. Colloquium Physical Measurements and Signatures in Remote Sensing*, Courchevel, France, Vol. **2**, pp. 501–504 (1991).
82. H. Y. Lee, "An application of NOAA AVHRR thermal data to the study of urban heat islands," *Atmos. Environ. Part B* **27**(1), 1–13 (1993).
83. J. Nichol, "Remote sensing of urban heat islands by day and night," *Photogramm. Eng. Remote Sens.* **71**(5), 613–621 (2005).
84. M. Tiangco, A. M. F. Lagmay, and J. Argete, "ASTER-based study of the night-time urban heat island effect in Metro Manila," *Int. J. Remote Sens.* **29**(10), 2799–2818 (2008).
85. L. Liu and Y. Zhang, "Urban heat island analysis using the Landsat TM data and ASTER data: a case study in Hong Kong," *Remote Sens.* **3**(7), 1535–1552 (2011).
86. K. Deilami et al., "Urban heat island effect: a systematic review of spatio-temporal factors, data, methods, and mitigation measures," *Int. J. Appl. Earth Obs. Geoinf.* **67**, 30–42 (2018).
87. E. Alves, "Seasonal and spatial variation of surface urban heat island intensity in a small urban agglomerate in Brazil," *Climate* **4**, 61 (2016).
88. Y. Makido et al., "Daytime variation of urban heat islands: the case study of Doha, Qatar," *Climate* **4**, 32 (2016).
89. N. T. Son et al., "Assessment of urbanization and urban heat islands in Ho Chi Minh City, Vietnam using Landsat data," *Sustainable Cities Soc.* **30**, 150–161 (2017).

90. H. Wang et al., "Surface urban heat island analysis of shanghai (China) based on the change of land use and land cover," *Sustainability* **9**(9), 1538 (2017).
91. W. Chen et al., "Evaluation of urbanization dynamics and its impacts on surface heat islands: a case study of Beijing, China," *Remote Sens.* **9**(5), 453 (2017).
92. A. Chen et al., "Effect of urban green patterns on surface urban cool islands and its seasonal variations," *Urban For. Urban Greening* **13**(4), 646–654 (2014).
93. U.S. Geological Survey, "August 22, 2013 - Landsat 8 Thermal Infrared Sensor (TIRS) Calibration Notice," <https://landsat.usgs.gov/august-22-2013-l8-tirs-calibration-notice> (July 2018).
94. A. Rasul et al., "A review on remote sensing of urban heat and cool islands," *Land* **6**(2), 38 (2017).
95. J. R. Irons, J. L. Dwyer, and J. A. Barsi, "The next Landsat satellite: the Landsat data continuity mission," *Remote Sens. Environ.* **122**, 11–21 (2012).
96. M. A. Wulder et al., "Opening the archive: how free data has enabled the science and monitoring promise of Landsat," *Remote Sens. Environ.* **122**, 2–10 (2012).
97. D. Zhou et al., "Climate-vegetation control on the diurnal and seasonal variations of surface urban heat islands in China," *Environ. Res. Lett.* **11**(7), 074009 (2016).
98. A. Mathew, S. Khandelwal, and N. Kaul, "Analysis of diurnal surface temperature variations for the assessment of surface urban heat island effect over Indian cities," *Energy Build.* **159**, 271–295 (2018).
99. R. Yao et al., "The influence of different data and method on estimating the surface urban heat island intensity," *Ecol. Indic.* **89**, 45–55 (2018).
100. J. Zhou et al., "Maximum nighttime urban heat island (UHI) intensity simulation by integrating remotely sensed data and meteorological observations," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **4**, 138–146 (2011).
101. J. Zhou et al., "Modelling the diurnal variations of urban heat islands with multi-source satellite data," *Int. J. Remote Sens.* **34**, 7568–7588 (2013).
102. H. Zhang et al., "Bibliometric analysis of global remote sensing research during 2010–2015," *ISPRS Int. J. Geo-Inf.* **6**(11), 332 (2017).
103. H. Bahi et al., "Effects of urbanization and seasonal cycle on the surface urban heat island patterns in the coastal growing cities: a case study of Casablanca, Morocco," *Remote Sens.* **8**(10), 829 (2016).
104. H. Du et al., "Influences of land cover types, meteorological conditions, anthropogenic heat and urban area on surface urban heat island in the Yangtze River Delta Urban Agglomeration," *Sci. Total Environ.* **571**, 461–470 (2016).
105. L. Gawuc and J. Struzewska, "Impact of MODIS quality control on temporally aggregated urban surface temperature and long-term surface urban heat island intensity," *Remote Sens.* **8**(5), 374 (2016).
106. M. Morabito et al., "The impact of built-up surfaces on land surface temperatures in Italian urban areas," *Sci. Total Environ.* **551–552**, 317–326 (2016).
107. C. Wang et al., "Spatio-temporal modeling of the urban heat island in the Phoenix metropolitan area: land use change implications," *Remote Sens.* **8**(3), 185 (2016).
108. S. Peng et al., "Surface urban heat island across 419 global big cities," *Environ. Sci. Technol.* **46**(2), 696–703 (2012).
109. S. Zhao, D. Zhou, and S. Liu, "Data concurrency is required for estimating urban heat island intensity," *Environ. Pollut.* **208**, 118–124 (2016).
110. D. Zhou et al., "Spatiotemporal trends of urban heat island effect along the urban development intensity gradient in China," *Sci. Total Environ.* **544**, 617–626 (2016).
111. N. Clinton and P. Gong, "MODIS detected surface urban heat islands and sinks: global locations and controls," *Remote Sens. Environ.* **134**, 294–304 (2013).
112. A. J. W. de Wit and C. A. van Diepen, "Crop growth modelling and crop yield forecasting using satellite-derived meteorological inputs," *Int. J. Appl. Earth Obs. Geoinf.* **10**(4), 414–425 (2008).
113. C. Daly, "Guidelines for assessing the suitability of spatial climate data sets," *Int. J. Climatol.* **26**(6), 707–721 (2006).

114. B. Christiansen, "Downward propagation and statistical forecast of the near-surface weather," *J. Geophys. Res. D* **110**(14), D14104 (2005).
115. E. N. Florio et al., "Integrating AVHRR satellite data and NOAA ground observations to predict surface air temperature: a statistical approach," *Int. J. Remote Sens.* **25**(15), 2979–2994 (2004).
116. K. Stahl et al., "Comparison of approaches for spatial interpolation of daily air temperature in a large region with complex topography and highly variable station density," *Agric. For. Meteorol.* **139**(3–4), 224–236 (2006).
117. J. V. Vogt, A. A. Viau, and F. Paquet, "Mapping regional air temperature fields using satellite-derived surface skin temperatures," *Int. J. Climatol.* **17**(14), 1559–1579 (1997).
118. Y. Chen et al., "Enhanced statistical estimation of air temperature incorporating nighttime light data," *Remote Sens.* **8**(8), 656 (2016).
119. Z. Sun et al., "Evaluation of the VI-Ts method for estimating the land surface moisture index and air temperature using ASTER and modis data in the North China Plain," *Int. J. Remote Sens.* **32**(22), 7257–7278 (2011).
120. C. Wloczyk et al., "Estimation of instantaneous air temperature above vegetation and soil surfaces from Landsat 7 ETM+ data in northern Germany," *Int. J. Remote Sens.* **32**(24), 9119–9136 (2011).
121. R. Pelta and A. A. Chudnovsky, "Spatiotemporal estimation of air temperature patterns at the street level using high resolution satellite imagery," *Sci. Total Environ.* **579**, 675–684 (2017).
122. S. Zhu et al., "Preliminary verification of instantaneous air temperature estimation for clear sky conditions based on SEBAL," *Meteorol. Atmos. Phys.* **129**(1), 71–81 (2017).
123. N. Janatian et al., "A statistical framework for estimating air temperature using MODIS land surface temperature data," *Int. J. Climatol.* **37**(3), 1181–1194 (2017).
124. J. Rhee and J. Im, "Estimating high spatial resolution air temperature for regions with limited *in situ* data using MODIS products," *Remote Sens.* **6**(8), 7360–7378 (2014).
125. L. Zeng et al., "Estimation of daily air temperature based on MODIS land surface temperature products over the Corn Belt in the US," *Remote Sens.* **7**(1), 951–970 (2015).
126. F. Huang et al., "Air temperature estimation with MODIS data over the Northern Tibetan Plateau," *Adv. Atmos. Sci.* **34**(5), 650–662 (2017).
127. C. Yoo et al., "Estimation of daily maximum and minimum air temperatures in urban landscapes using MODIS time series satellite data," *ISPRS J. Photogramm. Remote Sens.* **137**, 149–162 (2018).
128. S. Emamifar, A. Rahimikhoob, and A. A. Noroozi, "Daily mean air temperature estimation from MODIS land surface temperature products based on M5 model tree," *Int. J. Climatol.* **33**(15), 3174–3181 (2013).
129. Y. Xu, A. Knudby, and H. C. Ho, "Estimating daily maximum air temperature from MODIS in British Columbia, Canada," *Int. J. Remote Sens.* **35**(24), 8108–8121 (2014).
130. H. Meyer et al., "Mapping daily air temperature for Antarctica based on MODIS LST," *Remote Sens.* **8**(9), 732 (2016).
131. H. Zhang et al., "Estimating daily air temperatures over the Tibetan Plateau by dynamically integrating MODIS LST data," *J. Geophys. Res.* **121**(19), 11,425–11,441 (2016).
132. P. T. Noi, J. Degener, and M. Kappas, "Comparison of multiple linear regression, cubist regression, and random forest algorithms to estimate daily air surface temperature from dynamic combinations of MODIS LST data," *Remote Sens.* **9**, 398 (2017).
133. Y. Xu et al., "Mapping monthly air temperature in the Tibetan Plateau from MODIS data based on machine learning methods," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **11**(2), 345–354 (2018).
134. Y. Z. Yang, W. H. Cai, and J. Yang, "Evaluation of MODIS land surface temperature data to estimate near-surface air temperature in Northeast China," *Remote Sens.* **9**(5), 410 (2017).
135. P. T. Noi, M. Kappas, and J. Degener, "Estimating daily maximum and minimum land air surface temperature using MODIS land surface temperature data and ground truth data in Northern Vietnam," *Remote Sens.* **8**(12), 1002 (2016).

136. W. Zhou et al., "Estimating high resolution daily air temperature based on remote sensing products and climate reanalysis datasets over glacierized basins: a case study in the Langtang Valley, Nepal," *Remote Sens.* **9**(9), 959 (2017).
137. Z. Wan, "New refinements and validation of the MODIS land-surface temperature/emissivity products," *Remote Sens. Environ.* **112**, 59–74 (2008).
138. S. A. Ackerman et al., "Cloud detection with MODIS. Part II: validation," *J. Atmos. Oceanic Technol.* **25**, 1073–1086 (2008).
139. S. N. Williamson et al., "Evaluating cloud contamination of clear-sky MODIS Terra daytime land surface temperatures using ground-based meteorology station observations," *J. Clim.* **26**, 1551–1560 (2013).
140. J. Kim and T. S. Hogue, "Evaluation of a MODIS triangle-based evapotranspiration algorithm for semi-arid regions," *J. Appl. Remote Sens.* **7**(1), 073493 (2013).
141. D. A. Walker et al., "Environment, vegetation and greenness (NDVI) along the North America and Eurasia Arctic transects," *Environ. Res. Lett.* **7**(1), 015504 (2012).
142. S. B. Duan and Z. L. Li, "Spatial downscaling of MODIS land surface temperatures using geographically weighted regression: case study in Northern China," *IEEE Trans. Geosci. Remote Sens.* **54**, 6458–6469 (2016).
143. C. Zeng et al., "A two-step framework for reconstructing remotely sensed land surface temperatures contaminated by cloud," *ISPRS J. Photogramm. Remote Sens.* **141**, 30–45 (2018).
144. M. Jin and R. E. Dickinson, "A generalized algorithm for retrieving cloudy sky skin temperature from satellite thermal infrared radiances," *J. Geophys. Res. Atmos.* **105**(D22), 27037–27047 (2000).
145. L. Sun et al., "Reconstructing daily clear-sky land surface temperature for cloudy regions from MODIS data," *Comput. Geosci.* **105**, 10–20 (2017).
146. T. I. Østby, T. V. Schuler, and S. Westermann, "Severe cloud contamination of MODIS land surface temperatures over an Arctic ice cap, Svalbard," *Remote Sens. Environ.* **142**, 95–102 (2014).
147. D. Mutiibwa, S. Strachan, and T. Albright, "Land surface temperature and surface air temperature in complex terrain," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **8**(10), 4762–4774 (2015).
148. Y. Xu, Y. Shen, and Z. Wu, "Spatial and temporal variations of land surface temperature over the Tibetan Plateau based on harmonic analysis," *Mt. Res. Dev.* **33**(1), 85–94 (2013).
149. T. Shuai et al., "A spectral angle distance-weighting reconstruction method for filled pixels of the MODIS land surface temperature product," *IEEE Geosci. Remote Sens. Lett.* **11**(9), 1514–1518 (2014).
150. M. Neteler, "Estimating daily land surface temperatures in mountainous environments by reconstructed MODIS LST data," *Remote Sens.* **2**(1), 333–351 (2010).
151. B. Huang et al., "Generating high spatiotemporal resolution land surface temperature for urban heat island monitoring," *IEEE Geosci. Remote Sens. Lett.* **10**(5), 1011–1015 (2013).
152. K. Liu et al., "Quantifying spatial-temporal pattern of urban heat island in Beijing: an improved assessment using land surface temperature (LST) time series observations from LANDSAT, MODIS, and Chinese new satellite GaoFen-1," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **9**(5), 2028–2042 (2016).
153. S. Mukherjee, P. K. Joshi, and R. D. Garg, "Analysis of urban built-up areas and surface urban heat island using downscaled MODIS derived land surface temperature data," *Geocarto Int.* **32**(8), 900–918 (2017).
154. K. Hazaymeh and Q. K. Hassan, "Fusion of MODIS and Landsat-8 surface temperature images: a new approach," *PLoS One* **10**(3), e0117755 (2015).
155. R. R. Colditz et al., "TiSeG: a flexible software tool for time-series generation of MODIS data utilizing the quality assessment science data set," *IEEE Trans. Geosci. Remote Sens.* **46**(10), 3296–3308 (2008).

Thanh Noi Phan received his BS degree in geodesy from Hanoi University of Mining and Geology, Vietnam, in 2007, and his MSc degree in natural resources management from Asian Institute of Technology, Thailand, in 2012. Currently, he is pursuing his PhD from

the Georg-August University of Göttingen, Germany. His research interests include remote sensing of environment, image analysis, time-series analysis, vegetation monitoring and dynamics, and machine learning.

Martin Kappas is a professor of geography and head of the Department of Cartography, GIS and Remote Sensing, Georg-August University of Göttingen. He received his MSc degree in geography from the University of Bonn in 1989, and his PhD in environmental sciences/climatology from the University of Mannheim (“summa cum laude”) in 1993. His main research theme is the use of remote sensing and GIS for land use/cover change and climate change studies.