Diagnosis of malignant pleural mesothelioma from pleural fluid by Fourier transform-infrared spectroscopy coupled with chemometrics (Erratum)

Sherif Abbas
Nihal Simsek Ozek
Salih Emri
Deniz Koksall
Mete Severcan
Feride Severcana

Diagnosis of malignant pleural mesothelioma from pleural fluid by Fourier transform-infrared spectroscopy coupled with chemometrics (Erratum)

Sherif Abbas,^{a,b} Nihal Simsek Ozek,^{a,c,d} Salih Emri,^{e,f} Deniz Koksai,^e Mete Severcan,^g and Feride Severcana^{h,i}

^aMiddle East Technical University, Department of Biological Sciences, Ankara, Turkey

^bAin Shams University, Physics Department, Cairo, Egypt

^cAtaturk University, Department of Biology, Erzurum, Turkey

^dAtaturk University, East Anatolian High Technology Research and Application Center (DAYTAM), Erzurum, Turkey

^eHacettepe University, Department of Chest Diseases, Ankara, Turkey

^fAltinbas University, Department of Chest Diseases, Faculty of Medicine, Istanbul, Turkey

^gMiddle East Technical University, Department of Electrical and Electronic Engineering, Ankara, Turkey

^hAltinbas University, Department of Biophysics, Faculty of Medicine, Istanbul, Turkey

ⁱAltinbas University, Graduate Program of Biomedical Sciences, Institute of Health Sciences, Istanbul, Turkey

[DOI: [10.1117/1.JBO.23.10.109801](https://doi.org/10.1117/1.JBO.23.10.109801)]

This article [J. Biomed. Opt. 23(10), 105003(2018)] was originally published online on 13 October 2018 with an error on page 10, which misstated the average accuracy of test sample classification.

The original sentence states, “Cooman’s plot of MPM versus LC and MPM versus BPE models demonstrated the correct classification of tested samples from each group with an average of 95.6% accuracy.”

The corrected sentence states, “Cooman’s plot of MPM versus LC and MPM versus BPE models demonstrated the correct classification of tested samples from each group with an average of 96.2% accuracy.”

This article was corrected online on 16 October 2018. It appears correctly in print.